

For Research Use Only

SUPT16H Monoclonal antibody

Catalog Number: 68243-1-Ig **Featured Product**



Basic Information

Catalog Number: 68243-1-Ig	GenBank Accession Number: NM_007192	Purification Method: Protein G purification
Size: 150ul , Concentration: 1000 µg/ml by Nanodrop;	GeneID (NCBI): 11198	CloneNo.: 1E4C4
Source: Mouse	Full Name: suppressor of Ty 16 homolog (S. cerevisiae)	Recommended Dilutions: WB 1:5000-1:50000 IF 1:500-1:2000
Isotype: IgG1	Calculated MW: 120 kDa	
Immunogen Catalog Number: AG29459	Observed MW: 135 kDa	

Applications

Tested Applications:
IF, WB, ELISA

Species Specificity:
Human, mouse, rat

Positive Controls:
WB : LNCaP cells, MCF-7 cells, HeLa cells, HEK-293 cells, HepG2 cells, Jurkat cells, HSC-T6 cells, NIH/3T3 cells, RAW 164.7 cells

IF : HeLa cells,

Background Information

SUPT16H, also named as FACT140, FACTP140, SPT16 and CDC68, belongs to the peptidase M24 family and SPT16 subfamily. SUPT16H is a component of the FACT complex, a general chromatin factor that acts to reorganize nucleosomes. The FACT complex is involved in multiple processes that require DNA as a template such as mRNA elongation, DNA replication and DNA repair. During transcription elongation the FACT complex acts as a histone chaperone that both destabilizes and restores nucleosomal structure. It facilitates the passage of RNA polymerase II and transcription by promoting the dissociation of one histone H2A-H2B dimer from the nucleosome, then subsequently promotes the reestablishment of the nucleosome following the passage of RNA polymerase II. The FACT complex is probably also involved in phosphorylation of 'Ser-392' of p53/TP53 via its association with CK2 (casein kinase II). It also involved in vitamin D-coupled transcription regulation via its association with the WINAC complex, a chromatin-remodeling complex recruited by vitamin D receptor (VDR), which is required for the ligand-bound VDR-mediated transrepression of the CYP27B1 gene.

Storage

Storage:
Store at -20°C. Stable for one year after shipment.

Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



WB result of SUPT16H antibody (68243-1-Ig; 1:30000); incubated at room temperature for 1.5 hours) with sh-Control and sh-SUPT16H transfected MCF-7 cells.



Various lysates were subjected to SDS PAGE followed by western blot with 68243-1-Ig (SUPT16H antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Lamin B1 Monoclonal antibody (HRP-66095) as loading control.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using SUPT16H antibody (68243-1-Ig, Clone: 1E4C4) at dilution of 1:1000 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).