

For Research Use Only

PECR Monoclonal antibody

Catalog Number: 68248-1-Ig



Basic Information

Catalog Number: 68248-1-Ig	GenBank Accession Number: BC002529	Purification Method: Protein G purification
Size: 150ul , Concentration: 1000 µg/ml by Nanodrop;	GeneID (NCBI): 55825	CloneNo.: 2D3G1
Source: Mouse	Full Name: peroxisomal trans-2-enoyl-CoA reductase	Recommended Dilutions: WB 1:5000-1:50000
Isotype: IgG1	Calculated MW: 33 kDa	
Immunogen Catalog Number: AG7022	Observed MW: 33 kDa	

Applications

Tested Applications:

WB, ELISA

Species Specificity:

Human, Mouse, Rat

Positive Controls:

WB : LNCaP cells, HSC-T6 cells, HeLa cells, HepG2 cells,
Jurkat cells, NIH/3T3 cells

Background Information

Peroxisomal trans-2-enoyl-CoA reductase (PECR) is also called TERP and TECP, which is located on chromosome q35 with 86627 bases. It can participate in carbon chain elongation in fatty acid metabolism and catalyze the last reaction of four long chain fatty acid elongation cycles. Each cycle of PECR adds two carbons to the long chain and very long chain fatty acid (VLCFA) chains, reducing the intermediate of trans-2,3-enoyl coenzyme A fatty acid to acyl coenzyme A, which can further extend the carbon chain by entering a new elongation cycle. Meanwhile, PECR is a peroxisome protein involved in fatty acid synthesis and plays an important role in milk fat synthesis. The molecular mass of PECR is 33 kDa. (PMID: 31467878)

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68248-1-Ig (PECR antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Beta Actin Monoclonal antibody (HRP-66009) as loading control.