

For Research Use Only

POLR3D Monoclonal antibody

Catalog Number: 68263-1-Ig

Featured Product



Basic Information

Catalog Number:

68263-1-Ig

Size:

150ul , Concentration: 1000 ug/ml by Nanodrop;

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG7646

GenBank Accession Number:

BC004484

GeneID (NCBI):

661

UNIPROT ID:

P05423

Full Name:

polymerase (RNA) III (DNA directed) polypeptide D, 44kDa

Calculated MW:

44 kDa

Observed MW:

44 kDa

Purification Method:

Protein G purification

CloneNo.:

2F5G4

Recommended Dilutions:

WB 1:2000-1:10000

Applications

Tested Applications:

WB, FC (Intra), ELISA

Species Specificity:

human, rabbit

Positive Controls:

WB : LNCaP cells, rabbit brain tissue, K-562 cells, HEK-293 cells, HeLa cells, Jurkat cells

Background Information

DNA-dependent RNA polymerase catalyzes the transcription of DNA into RNA using the four ribonucleoside triphosphates as substrates. Polymerase (RNA) III (DNA directed) polypeptide D (POLR3D) plays a key role in sensing and limiting infection by intracellular bacteria and DNA viruses and acts as nuclear and cytosolic DNA sensor involved in innate immune response. POLR3D can sense non-self dsDNA that serves as template for transcription into dsRNA.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68263-1-Ig (POLR3D antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.



WB result of POLR3D antibody (68263-1-Ig; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-POLR3D transfected HEK-293 cells.



1x10⁶ HeLa cells were intracellularly stained with 0.8 ug POLR3D Monoclonal antibody (68263-1-Ig, Clone:2F5G4) and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1)(red), or 0.8 ug Mouse IgG1 isotype control Mouse McAb (66360-1-Ig, Clone: 1F8D3) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).