For Research Use Only

TMPRSS2 Monoclonal antibody

Catalog Number:68264-1-lg



| Basic Information | Catalog Number: 68264-1-lg | GenBank Accession Number: BC051839 | Purification Method: Protein G purification |
|------------------------|--|---|--|
| | Size: 150ul, Concentration: 1000 ug/ml by Nanodrop; Source: Mouse Isotype: IgG1 Immunogen Catalog Number: AG5824 | GenelD (NCBI): 7113 | CloneNo.: 1H7E9 |
| | | UNIPROT ID: | Recommended Dilutions: WB 1:1000-1:8000 |
| | | O15393 Full Name: | WB 1:1000-1:8000 |
| | | transmembrane protease, serine 2 | |
| | | Calculated MW: 54 kDa | |
| | | Observed MW: 50 kDa | |
| Applications | Tested Applications: | Positive Controls: | |
| | WB, ELISA Species Specificity: Human, Pig | WB : PC-3 cells, pig kidney tissue, LNCaP cells, Caco-2 cells, COLO 320 cells, HT-29 cells, human testis tissue | |
| | TMPRSS2, also named as PRSS10, is a type II transmembrane serine protease which is highly expressed by the epithelium of the human prostate gland. TMPRSS2 may contribute to prostate tumour metastasis via the activation of PAR-2. TMPRSS2 is a Serine protease that proteolytically cleaves and activates the viral spike glycoproteins which facilitate virus-cell membrane fusions. TMPRSS2 is a host cell factor that is critical for the spread of several clinically relevant viruses, including influenza A viruses and coronaviruses (PMID: 23468491, 30626688). SARS-CoV-2 uses the SARS-CoV receptor ACE2 for entry and the serine protease TMPRSS2 for S protein priming. The initial spike protein priming by TMPRSS2 is essential for the entry and viral spread of SARS-CoV-2 through interaction with the ACE2 receptor (PMID: 32142651, 30626688). Camostat mesylate, an inhibitor of TMPRSS2, can block SARS-CoV-2 infection of lung cells (PMID: 32142651). The MW of TMPRSS2 is about 65-70 kDa. It can be cleaved into some chains with MW 54 kDa, 31 kDa and 26 kDa (PMID: 25734995, 20382709, 26018085). | | |
| Background Information | of PAR-2. TMPRSS2 is a Serine proteas which facilitate virus-cell membrane clinically relevant viruses, including 2 uses the SARS-CoV receptor ACE2 fo spike protein priming by TMPRSS2 is the ACE2 receptor (PMID: 32142651, 3 2 infection of lung cells (PMID: 32142 | te that proteolytically cleaves and fusions. TMPRSS2 is a host cell fac influenza A viruses and coronaviru or entry and the serine protease TM essential for the entry and viral sp i0626688). Camostat mesylate, ar 651). The MW of TMPRSS2 is about | activates the viral spike glycoproteins tor that is critical for the spread of several ses (PMID: 23468491, 30626688). SARS-CoV- PRSS2 for S protein priming. The initial read of SARS-CoV-2 through interaction with inhibitor of TMPRSS2, can block SARS-CoV- 65-70 kDa. It can be cleaved into some |
| Storage | of PAR-2. TMPRSS2 is a Serine proteas which facilitate virus-cell membrane clinically relevant viruses, including 2 uses the SARS-CoV receptor ACE2 fo spike protein priming by TMPRSS2 is the ACE2 receptor (PMID: 32142651, 3 2 infection of lung cells (PMID: 32142 | e that proteolytically cleaves and fusions. TMPRSS2 is a host cell fac influenza A viruses and coronaviru or entry and the serine protease TM essential for the entry and viral sp i0626688). Camostat mesylate, ar 651). The MW of TMPRSS2 is about 6 kDa (PMID: 25734995, 20382709) er shipment. | activates the viral spike glycoproteins tor that is critical for the spread of several ses (PMID: 23468491, 30626688). SARS-CoV PRSS2 for S protein priming. The initial read of SARS-CoV-2 through interaction with inhibitor of TMPRSS2, can block SARS-CoV- 65-70 kDa. It can be cleaved into some |

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68264-1-lg (TMPRSS2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRPconjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.



pig kidney tissue were subjected to SDS PAGE followed by western blot with 68264-1-1g (TMPRSS2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 68264-1-lg (TMPRSS2 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.