

For Research Use Only

PDXDC1 Monoclonal antibody

Catalog Number: 68278-1-Ig



Basic Information

Catalog Number: 68278-1-Ig	GenBank Accession Number: BC033748	Purification Method: Protein A purification
Size: 150ul , Concentration: 1000 ug/ml by Nanodrop;	GeneID (NCBI): 23042	CloneNo.: 1E6E2
Source: Mouse	UNIPROT ID: Q6P996	Recommended Dilutions: WB 1:5000-1:50000 IF/ICC 1:400-1:1600
Isotype: IgG1	Full Name: pyridoxal-dependent decarboxylase domain containing 1	
Immunogen Catalog Number: AG15489	Calculated MW: 788 aa, 87 kDa	
	Observed MW: 87 kDa	

Applications

Tested Applications: WB, IF/ICC, ELISA	Positive Controls:
Species Specificity: Human, mouse, rat, pig	WB : LNCaP cells, NIH/3T3 cells, HEK-293 cells, Jurkat cells, K-562 cells, pig colon tissue, rat colon tissue IF/ICC : MCF-7 cells,

Background Information

PDXDC1 (pyridoxal-dependent decarboxylase domain containing 1) is a putative enzyme that could metabolize catecholamine neurotransmitters (PMID:28485732). With prior evidence for involvement with glioblastoma from other previously reported experimental settings, and contains the lead single nucleotide polymorphism (rs3198697) from the linkage analysis of the chromosome 16 region (PMID:32644145). Moreover, PDXDC1 is involved in the catalysis of the nonhydrolytic addition or removal of a carboxyl group to or from a compound (PMID:29277971).

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 68278-1-Ig (PDXDC1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control.



Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using PDXDC1 antibody (68278-1-Ig, Clone: 1E6E2) at dilution of 1:800 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red).



Immunofluorescent analysis of (-20°C Methanol) fixed MCF-7 cells using PDXDC1 antibody (68278-1-Ig, Clone: 1E6E2) at dilution of 1:800 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L).



NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 68278-1-Ig (PDXDC1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control.