## For Research Use Only

## MICAL1 Monoclonal antibody

Catalog Number: 68297-1-Ig



**Basic Information** 

Catalog Number: GenBank Accession Number:

68297-1-lg BC052983 GeneID (NCBI): Size:

150ul, Concentration: 1000 ug/ml by 64780 Nanodrop: **UNIPROT ID:** 

Q8TDZ2 Mouse Full Name:

Isotype: microtubule associated

lgG1 monoxygenase, calponin and LIM

domain containing 1 Immunogen Catalog Number: Calculated MW: AG6578

> 118 kDa Observed MW: 120 kDa

**Purification Method:** 

Protein G purification

CloneNo.: 3F5D4

Recommended Dilutions:

WB 1:5000-1:50000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IF/ICC 1:1000-1:4000

**Applications** 

**Tested Applications:** 

WB, IF/ICC, FC (Intra), IP, ELISA

Species Specificity: Human, mouse, rat

WB: A549 cells, RAW 264.7 cells, HSC-T6 cells, HeLa cells, HEK-293 cells, HepG2 cells, Jurkat cells, MOLT-4

cells, K-562 cells IP: HeLa cells. IF/ICC : HeLa cells.

**Background Information** 

MICALs (Molecules Interacting with CasL) are atypical multidomain flavoenzymes with diverse cellular functions. There are three known isoforms, MICAL1, MICAL2 and MICAL3, as well as the MICAL-like proteins MICAL-L1 and MICAL-L2. MICAL1 has four conserved domains: an N-terminal flavin adenine dinucleotide (FAD) binding domain, a calponin homology (CH) domain, a Lin11, Isl-1 and Mec-3 (LIM) domain and a C-terminal coiled-coil (CC) domain. MICAL1 is reported to regulate actin stress fibers and be required for normal actin organization. It may also be involved in apoptosis through binding with NDR (nuclear Dbf2-related) kinases. This antibody specially recognizes MICAL1.

Storage


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


\*\*\* 20ul sizes contain 0.1% BSA

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 68297-1-lg (MICAL1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with Lamin B1 Monoclonal antibody (66095-1-lg) as loading control.




IP result of anti-MICAL1 (IP:68297-1-Ig, 4ug; Detection:68297-1-Ig 1:2000) with HeLa cells lysate 1720 ug.




Immunofluorescent analysis of (4% PFA) fixed HeLa cells using MICAL1 antibody (68297-1-lg, Clone: 3F5D4) at dilution of 1:2000 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L), Alpha Tubulin antibody (11224-1-AP, red).



1X10^6 HeLa cells were intracellularly stained with 0.4 ug Anti-Human MICAL1 (68297-1-1g, Clone:3F5D4) and CoraLite® 488-Conjugated Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



HSC-T6 cells were subjected to SDS PAGE followed by western blot with 68297-1-1g (MICAL1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



RAW 264.7 cells were subjected to SDS PAGE followed by western blot with 68297-1-1g (MICAL1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.