

For Research Use Only

# m7G Monoclonal antibody

Catalog Number: 68302-1-Ig **1 Publications**



## Basic Information

|                 |  |                           |                         |
|-----------------|--|---------------------------|-------------------------|
| Catalog Number: | 68302-1-Ig                                     | GenBank Accession Number: | Purification Method:    |
| Size:           | 150ul , Concentration: 1000 ug/ml by Nanodrop; | GeneID (NCBI):            | Protein A purification  |
| Source:         | Mouse  | Full Name:                | CloneNo.: 2C3E10        |
| Isotype:        | IgG2b  |                           | Recommended Dilutions:  |
|                 |  |                           | IHC 1:1000-1:4000       |
|                 |  |                           | Dot Blot 1:2500-1:10000 |

## Applications

|  |   |
|--|---|
| Tested Applications:   | Positive Controls:  |
| IHC, Dot Blot, ELISA   | IHC : mouse testis tissue, human colon cancer tissue, human liver cancer tissue, human pancreas cancer tissue |
| Species Specificity:   | Dot Blot : HeLa cells,  |
| chemical compound  |   |
| Cited Species:   |   |
| mouse  |   |
| <b>Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0</b> |   |

## Background Information

7-Methylguanosine (m7G) is a modified purine nucleoside. It is a methylated version of guanosine and when found in human urine, it may be a biomarker of some types of cancer. In the RNAs, 7-methylguanosine have been used to study and examine the reaction evolving methylguanosine. It also plays a role in mRNA as a blocking group at its 5'-end. The m7G modification actively participates in biological and pathological functions by affecting the metabolism of various RNA molecules, including messenger RNA, ribosomal RNA, microRNA, and transfer RNA. Increasing evidence indicates a critical role for m7G in human disease development, especially cancer, and aberrant m7G levels are closely associated with tumorigenesis and progression via regulation of the expression of multiple oncogenes and tumor suppressor genes.

Protocol for Dot Blot:  
<https://www.ptglab.com/protocol/68302-1-IgDotBlot.pdf>

## Notable Publications

| Author       | Pubmed ID | Journal       | Application |
|--------------|-----------|---------------|-------------|
| Zhanzhi Meng | 38867322  | Eur J Med Res |             |

## Storage


Storage:  
Store at -20°C. Stable for one year after shipment.  
Storage Buffer:  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Indirect ELISA and competitive ELISA results show that this antibody is specific to m7G. Indirect ELISA (blue curve, refer to top X-right Y axis) was performed by coating BSA conjugated m7G at 10ng/well followed by blocking with 1% BSA. Serial diluted primary antibody was added to the plates and incubated at 37°C. HRP-goat anti-mouse was used for detection. Competitive ELISA was performed similarly except that different concentration of m7G or its structure analogue




Total RNA was isolated from HeLa cell line and was dotted to NC membrane at different amount as indicated above the dots. The membrane was blocked with BSA and blotted with m7G antibody 68302-1-Ig at 1:5000 followed by incubation of HRP-goat anti-mouse secondary antibody. Signal was developed by ECL substrate. A parallel dot blot was performed using Mouse IgG2b isotype control Monoclonal antibody 66360-3-Ig at the same dose.




Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 68302-1-Ig (m7G antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human pancreas cancer tissue slide using 68302-1-Ig (m7G antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 68302-1-Ig (m7G antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse testis tissue slide using 68302-1-Ig (m7G antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse testis tissue slide using 68302-1-Ig (m7G antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).