

For Research Use Only

Ataxin 2 Monoclonal antibody

Catalog Number: 68316-1-Ig



Basic Information

| | | |
|--|--|--|
| Catalog Number: 68316-1-Ig | GenBank Accession Number: BC114546 | Purification Method: Caprylic acid/ammonium sulfate precipitation |
| Size: 150ul, Concentration: 1000 µg/ml by Bradford method using BSA as the standard; | GeneID (NCBI): 6311 | CloneNo.: 1E8A2 |
| Source: Mouse | Full Name: ataxin 2 | Recommended Dilutions: WB 1:2000-1:10000 IHC 1:250-1:1000 IF 1:200-1:800 |
| Isotype: IgM | Calculated MW: 1313 aa, 140 kDa | |
| Immunogen Catalog Number: AG17402 | Observed MW: 140-150 kDa | |

Applications

| | |
|--|---|
| Tested Applications: IF, IHC, WB, ELISA | Positive Controls: |
| Species Specificity: Human, Mouse, Rat | WB: LNCaP cells, HeLa cells, HEK-293 cells, HepG2 cells, MCF-7 cells, Jurkat cells, HSC-T6 cells, NIH/3T3 cells, 4T1 cells |
| Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 | IHC: rat brain tissue, IF: rat brain tissue, |

Background Information

ATXN2 contains a repeat structure with 22 or 23 triplets coding for glutamine and the (CAG)₈CAA(CAG)₄CAA(CAG)₈ sequence; expansion of this domain to a size ≥34 triplets with a pure CAG sequence primarily causes autosomal dominant SCA2 [PMID:18418684], while ATXN2 expansions with CAA interruptions were observed as the cause of Levo-dopa responsive Parkinson's disease [PMID:10668726]. ATXN2 expansions associated with ALS were reported to be interrupted by at least one CAA triplet [PMID:21537950]

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 68316-1-Ig (Ataxin 2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded rat brain tissue slide using 68316-1-Ig (Ataxin 2 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded rat brain tissue slide using 68316-1-Ig (Ataxin 2 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed rat brain tissue using Ataxin 2 antibody (68316-1-Ig, Clone: 1E8A2) at dilution of 1:400 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed rat brain tissue using Ataxin 2 antibody (68316-1-Ig, Clone: 1E8A2) at dilution of 1:400 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).