For Research Use Only

CCDC94 Monoclonal antibody

Catalog Number:68324-1-lg Featured Product



Purification Method:

Basic Information

Catalog Number: GenBank Accession Number:

68324-1-Ig BC000561 Protein A purification
Size: GeneID (NCBI): CloneNo.:

Size: GenelD (NCBI): CloneNo 150ul , Concentration: 1000 ug/ml by 55702 1D1G9

 Nanodrop;
 UNIPROT ID:
 Recommended Dilutions:

 Source:
 Q9BW85
 WB 1:5000-1:50000

 Mouse
 Full Name:
 IF/ICC 1:750-1:3000

Isotype: coiled-coil domain containing 94

IgG2aCalculated MW:Immunogen Catalog Number:323 aa, 37 kDaAG13772Observed MW:

40 kDa

Applications

Tested Applications: Positive Controls:

WB, IF/ICC, ELISA

WB: A549 cells, HeLa cells, HEK-293 cells, HepG2 cells,
Species Specificity:

Jurkat cells, K-562 cells, HSC-T6 cells, NIH/3T3 cells,

human, mouse, rat 4T1 cells

IF/ICC: A431 cells,

Background Information

CCDC 94, also known as YJU2, is a splicing factor, predicted to be involved in RNA splicing and negative regulation of DNA damage response, signal transduction by p53 class mediator. Part of U2-type catalytic step 1 spliceosome. And that it is required for pre-mRNA splicing both in vivo and in vitro. Yju2 acts in concert with an unidentified heat-resistant factor(s) in an ATP-independent manner to promote the first catalytic reaction of pre-mRNA splicing after Prp2-mediated structural rearrangement of the spliceosome. (PMID: 17515604)

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 68324-1-lg (CCDC94 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.

WB result of CCDC94 antibody (68324-1-Ig; 1:10000; incubated at room temperature for 1.5 hours) with sh-Control and sh-CCDC94 transfected HeIa rells.



Immunofluorescent analysis of (4% PFA) fixed A431 cells using CCDC94 antibody (68324-1-1g, Clone: 1D1G9) at dilution of 1:1500 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (RGAM002), CL594-phalloidin (red).