

For Research Use Only

Histone H3 Monoclonal antibody

Catalog Number: 68345-1-Ig **10 Publications**



Basic Information

Catalog Number: 68345-1-Ig	GenBank Accession Number: BC066245	Purification Method: Protein A purification
Size: 150ul, Concentration: 1000 µg/ml by Nanodrop;	GeneID (NCBI): 8350	CloneNo.: 1A2A3
Source: Mouse	Full Name: histone cluster 1, H3a	Recommended Dilutions: WB 1:5000-1:50000 IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate IHC 1:500-1:2000 IF 1:500-1:2000
Isotype: IgG2b	Observed MW: 15 kDa	

Applications

Tested Applications:
FC, IF, IHC, IP, WB, ELISA

Cited Applications:
IF, WB

Species Specificity:
Human, mouse, rat, chicken, zebrafish, wheat, pig, rabbit, hamster, dog, canine

Cited Species:
human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: LNCaP cells, HEK-293 cells, Jurkat cells, K-562 cells, HSC-T6 cells, NIH/3T3 cells, MDCK cells, CHO cells, HeLa cells, chicken brain tissue, zebrafish, wheat whole plant

IP: HeLa cells,

IHC: mouse testis tissue,

IF: MCF-7 cells, A431 cells, HeLa cells

Background Information

Histones are small, highly basic proteins that consist of a globular domain with unstructured N- and C-terminal tails protruding from the main structure. Histone H3 is one of the five main histones that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. In addition to their role in DNA compartmentalization, histones also play crucial roles in various biologic processes, including gene expression and regulation, DNA repair, chromatin condensation, cell cycle progression, chromosome segregation, and apoptosis. The ability of histones to regulate chromatin dynamics primarily originates from various posttranslational modifications carried out by histone-modifying enzymes.

Notable Publications

Author	Pubmed ID	Journal	Application
Jordan Chang	37991367	J Virol	WB
Shengjie Tang	37949309	Brain Res	WB
Pan Lei	37834220	Int J Mol Sci	WB

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 68345-1-Ig (HIST1H3A antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 68345-1-Ig (Histone H3 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




IP result of anti-Histone H3 (IP:68345-1-Ig, 4ug; Detection:68345-1-Ig 1:2000) with HeLa cells lysate 1800 ug.




Immunohistochemical analysis of paraffin-embedded mouse testis tissue slide using 68345-1-Ig (HIST1H3A antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (4% PFA) fixed A431 cells using Histone H3 antibody (68345-1-Ig, Clone: 1A2A3) at dilution of 1:1000 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-phalloidin (red).




Immunofluorescent analysis of (4% PFA) fixed HeLa cells using Histone H3 antibody (68345-1-Ig, Clone: 1A2A3) at dilution of 1:1000 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-phalloidin (red).




Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using Histone H3 antibody (68345-1-Ig, Clone: 1A2A3) at dilution of 1:1000 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-phalloidin (red).




Immunofluorescent analysis of (4% PFA) fixed HeLa cells using Histone H3 antibody (68345-1-Ig, Clone: 1A2A3) at dilution of 1:400 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-phalloidin (red).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using Histone H3 antibody (68345-1-Ig, Clone: 1A2A3) at dilution of 1:1000 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), Alpha Tubulin antibody (11224-1-AP, red).



Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using Histone H3 antibody (68345-1-Ig, Clone: 1A2A3, labeled with CoraLite647-conjugated AffiniPure F(ab')₂ Fragment Goat Anti-Mouse IgG (H+L), SA00014-10, magenta) at dilution of 1:400 and CoraLite Plus 488-conjugated ZO-1 antibody (CL488-21773, green), CHCHD6 antibody (66597-1-Ig, Clone: 2A11E9, labeled with FlexAble CoraLite Plus 555 Antibody



1X10⁶ HepG2 cells were intracellularly stained with 0.4 ug Anti-Human Histone H3 (68345-1-Ig, Clone:1A2A3) and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG2b Isotype Control (MPC-11) (65128-1-Ig, Clone: MPC-11) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).