For Research Use Only

CEA Monoclonal antibody

Catalog Number: 68377-1-Ig



Basic Information

Catalog Number: GenBank Accession Number:

68377-1-lg BC034671 Protein G purification Size: Genel D (NCBI): CloneNo.:

150ul , Concentration: 1000 µg/ml by 1048 2B8H6

Nanodrop; Full Name: Recommended Dilutions:
Source: carcinoembryonic antigen-related WB 1:5000-1:50000
Mouse cell adhesion molecule 5 IHC 1:2500-1:10000

Isotype: Calculated MW: IgG1 77 kDa

Observed MW: 180-200 kDa

Applications

Tested Applications:

IHC, WB, ELISA
Species Specificity:

Human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HaCaT cells, human saliva, HT-29 cells, human

Purification Method:

rectum tissue

IHC: human colon cancer tissue,

Background Information

Carcinoembryonic antigen (CEA), also known as CEACAM5 or CD66e, is a cell surface glycoprotein belonging to the immunoglobulin superfamily, mainly serving as a cell adhesion molecule mediating intercellular contact by both homophilic and heterophilic binding (PMID: 21731662). CEA inhibits anoikis and plays a role in tumorigenesis and metastasis. CEA has been found to be overexpressed in a wide variety of human cancers, including colon, breast, and lung (PMID: 17167768). CEA is a tumor marker and is routinely exploited for diagnosis.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Aliquoting is unnecessary for -20°C storage


Storage Buffe


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

*** 20ul sizes contain 0.1% BSA


Selected Validation Data








Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 68377-1-Ig (CEACAM5 antibody) at dilution of 1:5000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 68377-1-1g (CEA antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 68377-1-1g (CEA antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.