For Research Use Only

CEA/CD66e Monoclonal antibody

Catalog Number: 68377-1-Ig



Basic Information

Catalog Number: GenBank Accession Number:

68377-1-lg BC034671 Protein G purification GeneID (NCBI): Size: CloneNo.:

150ul, Concentration: 1000 ug/ml by 1048 2B8H6

Nanodrop: **UNIPROT ID:** Recommended Dilutions: P06731 WB 1:5000-1:50000 Mouse IHC 1:2500-1:10000 Full Name:

Isotype: carcinoembryonic antigen-related

lgG1 cell adhesion molecule 5

> Calculated MW: 77 kDa Observed MW: 180-200 kDa

Applications

Tested Applications:

WB, IHC, ELISA

Species Specificity:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: HaCaT cells, human saliva, HT-29 cells, human

Purification Method:

rectum tissue

IHC: human colon cancer tissue,

Background Information

Carcinoembryonic antigen (CEA), also known as CEACAM5 or CD66e, is a cell surface glycoprotein belonging to the immunoglobulin superfamily, mainly serving as a cell adhesion molecule mediating intercellular contact by both homophilic and heterophilic binding (PMID: 21731662). CEA inhibits anoikis and plays a role in tumorigenesis and metastasis. CEA has been found to be overexpressed in a wide variety of human cancers, including colon, breast, and lung (PMID: 17167768). CEA is a tumor marker and is routinely exploited for diagnosis.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


*** 20ul sizes contain 0.1% BSA


in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data








Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 68377-1-Ig (CEACAM5 antibody) at dilution of 1:5000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 68377-1-1g (CEA antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 68377-1-1g (CEA antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.