

For Research Use Only

ATPBD4 Monoclonal antibody

Catalog Number: 68397-1-Ig



Basic Information

Catalog Number: 68397-1-Ig	GenBank Accession Number: BC008485	Purification Method: Protein G purification
Size: 150ul , Concentration: 1000 ug/ml by Nanodrop;	GeneID (NCBI): 89978	CloneNo.: 2B12E1
Source: Mouse	UNIPROT ID: Q7L8W6	Recommended Dilutions: WB 1:5000-1:50000
Isotype: IgG1	Full Name: ATP binding domain 4	
Immunogen Catalog Number: AG33218	Calculated MW: 267 aa, 30 kDa	
	Observed MW: 28-30 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : A549 cells, K-562 cells, LNCaP cells, HeLa cells, HEK-293 cells
Species Specificity: Human	

Background Information

Diphthamide is a highly modified histidine in the transcription factor EEF2 that is conserved from yeast to human. ATPBD4, also known as DPH6, belongs to the Diphthine--ammonia ligase family. DPH6 amidase may catalyze the last step of diphthamide biosynthesis using ammonium and ATP to play a role in catalyzing the final step of diphthamide biosynthesis, amidation of diphthine to diphthamide (PubMed:23169644).

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68397-1-Ig (ATPBD4 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control.



K-562 cells were subjected to SDS PAGE followed by western blot with 68397-1-Ig (ATPBD4 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.