For Research Use Only

## ISG15 Monoclonal antibody

Catalog Number:68408-1-lg 1 Publications




Basic Information	Catalog Number: 68408-1-lg	GenBank Accession Num BC009507		Purification Method: Protein A purification	
	Size:	GenelD (NCBI):		CloneNo.:	
	150ul , Concentration: 1000 ug/ml by			3G9A2	
	Nanodrop;	UNIPROT ID:		Recommended Dilutions:	
	Source:	P05161		WB 1:2000-1:10000	
	Mouse	Full Name:			
	Isotype:	ISG15 ubiquitin-like modifier Calculated MW:			
	lgG2a				
	Immunogen Catalog Number: AG8953	165 aa, 18 kDa			
		Observed MW:			
		13-17 kDa			
Applications	Tested Applications:	Positive Controls:			
	WB, ELISA	W	WB : MDA-MB-231 cells, LNCaP cells, A431 cells, HeLa		
	Cited Applications: IHC	cells, HCT 116 cells, HeLa cells, DU 145 cells, Jurka cells			
	Species Specificity: Human				
	Cited Species:				
	human				
Background Information	alpha (IFNA) and -beta (IFNB). ISG15 ISGylation, which in mammals is kno like (UBL) domains and a highly cons conjugation motif. Intracellular ISG1	forms covalent conjugates wn to play a role in antivi erved C-terminal LRGG sec 5 are conjugated, via the L ubiquitination, the process	s with its targe ral immunity quence, the la RGG motif, to s of formation	ISG15 proteins possess two ubiquitin tter being known as the ubiquitin target proteins through a process calle of ubiquitin conjugates. Unconjugatec	
	alpha (IFNA) and -beta (IFNB). ISG15 ISGylation, which in mammals is kno like (UBL) domains and a highly const conjugation motif. Intracellular ISG19 ISGylation, which resembles largely i extracellular ISG15, which are release cytokine-like activity.	forms covalent conjugates wn to play a role in antivi erved C-terminal LRGG sec 5 are conjugated, via the L ubiquitination, the process	s with its targe ral immunity quence, the la RGG motif, to s of formation	et proteins in a process called . ISG15 proteins possess two ubiquitin- tter being known as the ubiquitin target proteins through a process calle of ubiquitin conjugates. Unconjugated rine cells, are known to possess	
Background Information Notable Publications	alpha (IFNA) and -beta (IFNB). ISG15 ISGylation, which in mammals is kno like (UBL) domains and a highly const conjugation motif. Intracellular ISG15 ISGylation, which resembles largely extracellular ISG15, which are release cytokine-like activity.	forms covalent conjugates wn to play a role in antivi erved C-terminal LRGG sec 5 are conjugated, via the L ubiquitination, the process ed from several types of h med ID Journal	s with its targe ral immunity quence, the la RGG motif, to s of formation uman and mu	et proteins in a process called .ISG15 proteins possess two ubiquitin- tter being known as the ubiquitin target proteins through a process calle of ubiquitin conjugates. Unconjugated rine cells, are known to possess Application	
	alpha (IFNA) and -beta (IFNB). ISG15 ISGylation, which in mammals is kno like (UBL) domains and a highly const conjugation motif. Intracellular ISG15 ISGylation, which resembles largely i extracellular ISG15, which are release cytokine-like activity.	forms covalent conjugates wn to play a role in antivi erved C-terminal LRGG sec 5 are conjugated, via the L ubiquitination, the process ed from several types of h	s with its targe ral immunity quence, the la RGG motif, to s of formation uman and mu	et proteins in a process called . ISG15 proteins possess two ubiquitin tter being known as the ubiquitin target proteins through a process calle of ubiquitin conjugates. Unconjugated rine cells, are known to possess	
Notable Publications	alpha (IFNA) and -beta (IFNB). ISG15 i ISGylation, which in mammals is kno like (UBL) domains and a highly const conjugation motif. Intracellular ISG15 ISGylation, which resembles largely i extracellular ISG15, which are release cytokine-like activity. Author Pub Haruto Nishida 390 Storage:	forms covalent conjugates wn to play a role in antivi erved C-terminal LRGG sec 5 are conjugated, via the L ubiquitination, the process ed from several types of h med ID Journal 153305 Exp Mol I	s with its targe ral immunity quence, the la RGG motif, to s of formation uman and mu	et proteins in a process called .ISG15 proteins possess two ubiquitin- tter being known as the ubiquitin target proteins through a process calle of ubiquitin conjugates. Unconjugated rine cells, are known to possess	
Notable Publications	alpha (IFNA) and -beta (IFNB). ISG15 ti ISGylation, which in mammals is kno like (UBL) domains and a highly const conjugation motif. Intracellular ISG15 ISGylation, which resembles largely is extracellular ISG15, which are release cytokine-like activity. Author Pub Haruto Nishida 390 Storage: Storage:	forms covalent conjugates wn to play a role in antivi erved C-terminal LRGG sec 5 are conjugated, via the L ubiquitination, the process ed from several types of h med ID Journal 153305 Exp Mol I	s with its targe ral immunity quence, the la RGG motif, to s of formation uman and mu	et proteins in a process called .ISG15 proteins possess two ubiquitin- tter being known as the ubiquitin target proteins through a process calle of ubiquitin conjugates. Unconjugated rine cells, are known to possess Application	
Notable Publications	alpha (IFNA) and -beta (IFNB). ISG15 ti ISGylation, which in mammals is kno like (UBL) domains and a highly const conjugation motif. Intracellular ISG19 ISGylation, which resembles largely to extracellular ISG15, which are release cytokine-like activity. Author Pub Haruto Nishida 390 Storage: Store at -20°C. Stable for one year after Storage Buffer:	forms covalent conjugates wn to play a role in antivi erved C-terminal LRGG sec 5 are conjugated, via the L ubiquitination, the process ed from several types of h med ID Journal 153305 Exp Mol I er shipment.	s with its targe ral immunity quence, the la RGG motif, to s of formation uman and mu	et proteins in a process called .ISG15 proteins possess two ubiquitin- tter being known as the ubiquitin target proteins through a process calle of ubiquitin conjugates. Unconjugated rine cells, are known to possess	
Notable Publications	alpha (IFNA) and -beta (IFNB). ISG15 ti ISGylation, which in mammals is kno like (UBL) domains and a highly const conjugation motif. Intracellular ISG15 ISGylation, which resembles largely is extracellular ISG15, which are release cytokine-like activity. Author Pub Haruto Nishida 390 Storage: Store at -20°C. Stable for one year after Storage Buffer: PBS with 0.02% sodium azide and 50°	forms covalent conjugates wn to play a role in antivi erved C-terminal LRGG sec 5 are conjugated, via the L ubiquitination, the process ed from several types of h med ID Journal 153305 Exp Mol I er shipment.	s with its targe ral immunity quence, the la RGG motif, to s of formation uman and mu	et proteins in a process called .ISG15 proteins possess two ubiquitin- tter being known as the ubiquitin target proteins through a process calle of ubiquitin conjugates. Unconjugated rine cells, are known to possess	
	alpha (IFNA) and -beta (IFNB). ISG15 ti ISGylation, which in mammals is kno like (UBL) domains and a highly const conjugation motif. Intracellular ISG19 ISGylation, which resembles largely to extracellular ISG15, which are release cytokine-like activity. Author Pub Haruto Nishida 390 Storage: Store at -20°C. Stable for one year after Storage Buffer:	forms covalent conjugates wn to play a role in antivi erved C-terminal LRGG sec 5 are conjugated, via the L ubiquitination, the process ed from several types of h med ID Journal 153305 Exp Mol I er shipment.	s with its targe ral immunity quence, the la RGG motif, to s of formation uman and mu	et proteins in a process called .ISG15 proteins possess two ubiquitin- tter being known as the ubiquitin target proteins through a process calle of ubiquitin conjugates. Unconjugated rine cells, are known to possess Application	

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68408-1-1g (ISG15 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.