

For Research Use Only

AQP4 Monoclonal antibody

Catalog Number: 68448-1-Ig



Basic Information

Catalog Number: 68448-1-Ig	GenBank Accession Number: BC022286	Purification Method: Protein A purification
Size: 150ul , Concentration: 1000 µg/ml by Nanodrop;	GeneID (NCBI): 361	CloneNo.: 3G7D3
Source: Mouse	Full Name: aquaporin 4	Recommended Dilutions: WB 1:5000-1:50000
Isotype: IgG2b	Calculated MW: 323 aa, 35 kDa	
Immunogen Catalog Number: AG9830	Observed MW: 35-37 kDa	

Applications

Tested Applications:

WB, ELISA

Species Specificity:

Human, mouse, rat, rabbit, pig

Positive Controls:

WB : pig brain tissue, rat spinal cord tissue, rabbit brain tissue, rat brain tissue, mouse brain tissue

Background Information

Aquaporins are specialized water transport channels in plasma membranes of water-permeable tissues. Aquaporin-4 (AQP4) is the most abundant water channel in the human central nervous system and is important to fluid movements in brain. Aquaporin-4 exists as two isoforms, a long (M1) isoform with translation initiation at Met-1, and a shorter (M23) isoform with translation initiation at Met-23, with molecular weights around 35-37 kDa and 32-34 kDa, respectively.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68448-1-Ig (AQP4 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



rat spinal cord tissue were subjected to SDS PAGE followed by western blot with 68448-1-Ig (AQP4 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.