For Research Use Only

GIMAP7 Monoclonal antibody

Catalog Number:68451-1-lg





Basic Information	Catalog Number: 68451-1-lg	GenBank Accession Number: BC027613	Purification Method: Protein G purification					
	Size: 150ul , Concentration: 1000 ug/ml by Nanodrop; Source: Mouse Isotype: IgG1 Immunogen Catalog Number: AG10740	GeneID (NCBI): / 168537 UNIPROT ID: Q8NHV1 Full Name: GTPase, IMAP family member 7 Calculated MW: 300 aa, 35 kDa Observed MW: 34 kDa	CloneNo.: 3E4A6 Recommended Dilutions: WB 1:5000-1:50000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IF/ICC 1:400-1:1600					
				Applications	Tested Applications: WB, IF/ICC, IP, ELISA		Positive Controls: WB : NK-92 cells, Jurkat cells, MOLT-4 cells, HUVEC cells IP : Jurkat cells,	
					Species Specificity: Human	cells		
						IF/ICC : He	oG2 cells,	
				Background Information	GIMAP7, also known as hIAN7 or IAN apparatus. GIMAP7 can promote oxid	•	ed in the endoplasmic reticulum and Golફ	
	syndrome by inhibiting the SHH signa antibody specially recognises the 34	alling pathway(PMID: 36581994). Th	n granulosa cells in Polycystic ovary ne MW of this protein is 34 kDa, and this					
Storage	, , , ,	alling pathway(PMID: 36581994). Th kDa protein. er shipment.						

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





IP result of anti-GIMAP7 (IP:68451-1-lg, 4ug; Detection:68451-1-lg 1:2000) with Jurkat cells lysate 1560 ug.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using GIMAP7 antibody (68451-1-lg, Clone: 3E4A6) at dilution of 1:800 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L).

Various lysates were subjected to SDS PAGE followed by western blot with 68451-1-lg (GIMAP7 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.