

For Research Use Only

NUDT10 Monoclonal antibody

Catalog Number: 68478-1-Ig



Basic Information

Catalog Number: 68478-1-Ig	GenBank Accession Number: BC050700	Purification Method: Protein A purification
Size: 150ul , Concentration: 1000 ug/ml by Nanodrop;	GeneID (NCBI): 170685	CloneNo.: 3A8D5
Source: Mouse	UNIPROT ID: Q8NFP7	Recommended Dilutions: WB 1:5000-1:50000
Isotype: IgG2b	Full Name: nudix (nucleoside diphosphate linked moiety X)-type motif 10	
Immunogen Catalog Number: AG11878	Calculated MW: 164 aa, 19 kDa	
	Observed MW: 20 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : U2OS cells, rat brain tissue, HEK-293 cells, rat testis tissue, mouse testis tissue, mouse brain tissue
Species Specificity: Human, mouse, rat	

Background Information

NUDT10 (nucleoside diphosphate linked moiety X-type motif 10), also named as DIPP3A or APS2, belongs to the subgroup of phosphohydrolases that preferentially attack diphosphoinositol polyphosphates (PMID: 12105228, 12689335). Five human isoforms of DIPP have been found and named as DIPP type 1 (NUDT3), type 2 α / β (NUDT4), type 3 α (NUDT10) and type 3 β (NUDT11), which have the similar molecular weight and high homology (PMID: 12105228).

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68478-1-Ig (NUDT10 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 68478-1-Ig (NUDT10 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.