For Research Use Only

Lactotransferrin/LTF Monoclonal antibody

Catalog Number: 68509-1-Ig



Purification Method:

CloneNo.:

3D1G3

Protein A purification

Recommended Dilutions:

WB 1:5000-1:50000

Basic Information

Catalog Number: GenBank Accession Number:

68509-1-lg BC015822
Size: GeneID (NCBI):

150ul , Concentration: 1000 ug/ml by 4057
Nanodrop; UNIPROT ID:
Source: P02788
Mouse Full Name:
Isotype: lactotransferrin
IgG2a Calculated MW:
Immunogen Catalog Number: 80 kDa
AG33645 Observed MW:

le: IHC 1:2000-1:8000

Applications

Tested Applications:

WB, IHC, ELISA

WB: human saliva,
Species Specificity:

human human HHC : human skin cancer tissue,

78 kDa

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

LTF(Lactotransferrin) is also named as GIG12, LF and belongs to the transferrin family. It is an iron binding transport protein which can bind two atoms of ferric iron in association with the binding of an anion, usually bicarbonate. LTF is also an important protein component of the innate immune system that is broadly distributed within the body fluids. It can contribute to the activation of both the innate and adaptive immune responses by promoting the recruitment of leukocytes and activation of dendritic cells(PMID:18453607). This protein has 2 isoforms produced by alternative promoter usage. The human LTF can be isolated two fragments, an N-terminal tryptic fragment having an MW of 30 kDa and a C-terminal tryptic fragment having an MW of 50 kDa by a mild tryptic digest(PMID:3790094).

Storage

Storage:


Store at -20°C. Stable for one year after shipment. Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



human saliva were subjected to SDS PAGE followed by western blot with 68509-1-lg (Lactoferrin antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded skin cancer slide using 68509-1-1g (LTF antibody) at dilution of 1:4000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).