

For Research Use Only

RNF8 Monoclonal antibody

Catalog Number: 68521-1-Ig



Basic Information

| | | |
|--|--|---|
| Catalog Number: 68521-1-Ig | GenBank Accession Number: BC007517 | Purification Method: Protein G purification |
| Size: 150ul , Concentration: 1000 ug/ml by Nanodrop; | GeneID (NCBI): 9025 | CloneNo.: 3G3H5 |
| Source: Mouse | UNIPROT ID: O76064 | Recommended Dilutions: WB 1:1000-1:4000 |
| Isotype: IgG1 | Full Name: ring finger protein 8 | |
| Immunogen Catalog Number: AG5292 | Calculated MW: 485 aa, 56 kDa | |
| | Observed MW: 65-70 kDa | |

Applications

| | |
|---|---|
| Tested Applications: WB, ELISA | Positive Controls: WB : HeLa cells, HEK-293 cells, K-562 cells, pig brain tissue, rabbit brain tissue, rat brain tissue, mouse brain tissue |
| Species Specificity: Human, mouse, rat, rabbit, pig | |

Background Information

RNF8, also named as KIAA0646 or E3 ubiquitin-protein ligase RNF8, is a 485 amino acid protein, which contains one RING-type zinc finger and one FHA domain. RNF8 localizes in the nucleus and belongs to the RNF8 family. E2-E3 ubiquitin ligase complex is composed of the RNF8 homodimer and a E2 heterodimer of UBE2N and UBE2V2. E3 ubiquitin-protein ligase that plays a key role in DNA damage signaling via 2 distinct roles: by mediating the 'Lys-63'-linked ubiquitination of histones H2A and H2AX and promoting the recruitment of DNA repair proteins at double-strand breaks (DSBs) sites, and by catalyzing 'Lys-48'-linked ubiquitination to remove target proteins from DNA damage sites.

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68521-1-Ig (RNF8 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.