

For Research Use Only

# MORC2 Monoclonal antibody, PBS Only



Catalog Number: 68530-1-PBS

## Basic Information

|   |  |   |
|---|--|---|
| <b>Catalog Number:</b><br>68530-1-PBS                       | <b>GenBank Accession Number:</b><br>BC136782           | <b>Purification Method:</b><br>Protein G purification |
| <b>Size:</b><br>100ug , Concentration: 1 mg/ml by Nanodrop; | <b>GeneID (NCBI):</b><br>22880                         | <b>CloneNo.:</b><br>1G12E7                            |
| <b>Source:</b><br>Mouse                                     | <b>UNIPROT ID:</b><br>Q9Y6X9                           |   |
| <b>Isotype:</b><br>IgG1                                     | <b>Full Name:</b><br>MORC family CW-type zinc finger 2 |   |
| <b>Immunogen Catalog Number:</b><br>AG19605                 | <b>Calculated MW:</b><br>970 aa, 111 kDa               |   |
|   | <b>Observed MW:</b><br>118 kDa                         |   |

## Applications

**Tested Applications:**  
WB, Indirect ELISA

**Species Specificity:**  
Human

## Background Information

MORC2 is essential for epigenetic silencing by the HUSH (human silencing hub) complex. It can be recruited by HUSH to target site in heterochromatin. MORC2 also recruits histone deacetylases, such as HDAC4, to promoter regions, causing local histone H3 deacetylation and transcriptional repression of genes.

## Storage


**Storage:**  
Store at -80°C.

**Storage Buffer:**  
PBS Only

For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68530-1-Ig (MORC2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 68530-1-PBS in a different storage buffer formulation.