

For Research Use Only

PUF60 Monoclonal antibody

Catalog Number: 68533-1-Ig **1 Publications**



Basic Information

Catalog Number: 68533-1-Ig	GenBank Accession Number: BC008875	Purification Method: Protein G purification
Size: 150ul , Concentration: 1000 ug/ml by Nanodrop;	GeneID (NCBI): 22827	CloneNo.: 1A3A1
Source: Mouse	UNIPROT ID: Q9UHX1	Recommended Dilutions: WB 1:5000-1:50000
Isotype: IgG1	Full Name: poly-U binding splicing factor 60kDa	
Immunogen Catalog Number: AG33930	Calculated MW: 60 kDa	
	Observed MW: 60 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : MCF-7 cells, mouse testis tissue
Cited Applications: WB	
Species Specificity: Human, mouse, rat, rabbit	
Cited Species: mouse	

Background Information

Poly-U-binding factor 60(PUF60) is a splicing factor homologous to and complementary in function to U2AF, which involves in pre-mRNA splicing, a stepwise process initiated by binding. Also PUF60 involves in other nuclear processes such as apoptosis and transcription regulation. It can interact with and repress FUBP1-induced transcriptional activation via the core-TFIIH basal transcription factor. However, it has no repression activity in xeroderma pigmentosum complementation group B (XPB) cells.

Notable Publications

Author	Pubmed ID	Journal	Application
Jingyi Ren	39282919	Food Funct	WB

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68533-1-Ig (PUF60 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



mouse testis tissue were subjected to SDS PAGE followed by western blot with 68533-1-Ig (PUF60 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.