

For Research Use Only

EXO1 Monoclonal antibody

Catalog Number: 68536-1-Ig



Basic Information

Catalog Number: 68536-1-Ig	GenBank Accession Number: BC007491	Purification Method: Protein G purification
Size: 150ul , Concentration: 1000 µg/ml by Nanodrop;	GeneID (NCBI): 9156	CloneNo.: 2D11D1
Source: Mouse	Full Name: exonuclease 1	Recommended Dilutions: WB 1:5000-1:50000
Isotype: IgG1	Calculated MW: 94 kDa	
Immunogen Catalog Number: AG9253	Observed MW: 115-120 kDa	

Applications

Tested Applications:
WB, ELISA

Species Specificity:
Human

Positive Controls:

WB : SH-SY5Y cells, HeLa cells, HEK-293T cells, MOLT-4 cells, TF-1 cells, K-562 cells

Background Information

EXO1, also named as HEX1, is a 846 amino acid protein, which belongs to the XPG/RAD2 endonuclease family. EXO1 subfamily. EXO1 is highly expressed in bone marrow, testis and thymus and at lower levels in colon, lymph nodes, ovary, placenta, prostate, small intestine, spleen and stomach. EXO1 as 5'→3' double-stranded DNA exonuclease, which may also possess a cryptic 3'→5' double-stranded DNA exonuclease activity. EXO1 functions in DNA mismatch repair (MMR) to excise mismatch-containing DNA tracts directed by strand breaks located either 5' or 3' to the mismatch. The calculated molecular weight of EXO1 is 94 kDa, but the modified EXO1 protein is about 115 kDa.

Storage

Storage:
Store at -20°C. Stable for one year after shipment.

Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68536-1-Ig (EXO1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.