## For Research Use Only

## GABPA Monoclonal antibody

Catalog Number: 68645-1-Ig



**Purification Method:** 

Protein A purification

CloneNo.:

**Basic Information** 

Catalog Number: GenBank Accession Number:

68645-1-lg BC035031 GeneID (NCBI): Size:

150ul, Concentration: 1000 ug/ml by 2551 2A7E3 Nanodrop: **UNIPROT ID:** Recommended Dilutions: Q06546 WB 1:2000-1:10000 Mouse IF/ICC 1:1000-1:4000

Full Name:

Isotype: GA binding protein transcription factor, alpha subunit 60kDa IgG2a

Immunogen Catalog Number: Calculated MW: AG33705 454 aa. 51 kDa

Observed MW: 56-60 kDa

**Applications** 

**Tested Applications:** Positive Controls:

WB, IF/ICC, ELISA WB: HeLa cells, HEK-293 cells, HepG2 cells, Jurkat Species Specificity: cells, K-562 cells, HSC-T6 cells, NIH/3T3 cells

human, mouse, rat IF/ICC: HEK-293 cells,

## **Background Information**

GA-binding protein alpha chain (GABP alpha subunit, GABPA, nuclear respiratory factor 2 subunit alpha, transcription factor E4TF 1-60) is one of three GA-binding protein transcription factor subunits which functions as a DNA-binding subunit.GABPA is a member of Ets family, binds to the Yap promoter and activates YAP transcription(23684612). Since this subunit shares identity with a subunit encoding the nuclear respiratory factor 2 gene, it is likely involved in activation of cytochrome oxidase expression and nuclear control of mitochondrial function. This subunit also shares identity with a subunit constituting the transcription factor E4TF1, responsible for expression of the adenovirus E4 gene. Because of its chromosomal localization and ability to form heterodimers with other polypeptides, this gene may play a role in the Down Syndrome phenotype.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68645-1-1g (GABPA antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HEK-293 cells using GABPA antibody (68645-1-1g, Clone: 2A7E3) at dilution of 1:2000 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (RGAM002), CL594-phalloidin (red).