For Research Use Only

ICAM-1/CD54 Monoclonal antibody, PBS Only (Detector)

www.ptglab.com

Catalog Number: 68693-1-PBS

Basic Information

Catalog Number:

GenBank Accession Number: BC015969

Purification Method: Protein G purification

68693-1-PBS

GeneID (NCBI):

Size:

100ug, Concentration: 1 mg/ml by

CloneNo.: 2C9G8

Nanodrop: Mouse

ENSEMBL Gene ID: ENSG00000090339 **UNIPROT ID:** P05362

Isotype: lgG1 Full Name: Immunogen Catalog Number:

intercellular adhesion molecule 1

EG0570

Calculated MW: 90 kDa

Applications

Tested Applications:

Sandwich ELISA, Indirect ELISA, Sample test

Species Specificity:

human

Product Information

68693-1-PBS targets ICAM-1/CD54 as part of a matched antibody pair:

MP50085-1: 65075-1-PBS capture and 68693-1-PBS detection (validated in Sandwich ELISA)

Unconjugated mouse monoclonal antibody pair in PBS only (BSA and azide free) storage buffer at a concentration of 1 mg/mL, ready for conjugation.

This conjugation ready format makes antibodies ideal for use in many applications including: ELISAs, multiplex assays requiring matched pairs, mass cytometry, and multiplex imaging applications. Antibody use should be optimized by the end user for each application and assay.

Storage


Storage: Store at -80°C.


Storage Buffer:


PBS only, pH7.3

in USA), or 1(312) 455-8498 (outside USA)


Selected Validation Data









Human peripheral blood mononuclear cells (PBMC) were cultured unstimulated or stimulated with 10 µg/mL PHA for 5 days. The mean ICAM-1/CD54 concentration was determined to be 2,139.3 pg/mL in unstimulated PBMC supernatant, 30,678.8 pg/mL in PHA stimulated PBMC supernatant.



Saliva of seven individual healthy human donors was measured. The human ICAM-1/CD54 concentration of detected samples was determined to be 5,597.4 pg/mL with a range of 2,577.6 - 11,698.1 pg/mL



Urine of seven individual healthy human donors was measured. The human ICAM-1/CD54 concentration of detected samples was determined to be 1,031.9 pg/mL with a range of 49.7 - 3,299.7 pg/mL



Serum of sixteen individual healthy human donors was measured. The human ICAM-1/CD54 concentration of detected samples was determined to be 260.08 ng/mL with a range of 142.73 - 650.42 ng/mL