

For Research Use Only

NPTX2 Monoclonal antibody

Catalog Number: 68709-1-Ig



Basic Information

Catalog Number: 68709-1-Ig	GenBank Accession Number: BC009924	Purification Method: Protein G purification
Size: 150ul , Concentration: 1000 µg/ml by Nanodrop;	GeneID (NCBI): 4885	CloneNo.: 2H8D8
Source: Mouse	UNIPROT ID: P47972	Recommended Dilutions: WB 1:5000-1:50000
Isotype: IgG1	Full Name: neuronal pentraxin II	
Immunogen Catalog Number: AG33808	Calculated MW: 47 kDa	
	Observed MW: 50-55 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : pig pituitary gland tissue, pig brain tissue, rabbit brain tissue, rat brain tissue, mouse brain tissue
Species Specificity: Human, mouse, rat, pig, rabbit	

Background Information

Neuronal pentraxins constitute a family of proteins that are homologous to C-reactive protein (CRP) and serum amyloid P component (SAP), including NPTX1, NPTX2, and the neuronal pentraxin receptor (NPTXR). NPTX2, also known as NARP (neuronal activity-regulated pentaxin), is a secreted protein involved in excitatory synapse formation. It also plays a role in the clustering of alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA)-type glutamate receptors at established synapses, resulting in non-apoptotic cell death of dopaminergic nerve cells. NPTX2 is highly up-regulated in Parkinson's disease (PD), suggesting it may be involved in the pathology of PD. (PMID: 12895424; 17987278; 10748068)

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68709-1-Ig (NPTX2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.