

For Research Use Only

KIAA0319L Monoclonal antibody

Catalog Number: 68717-1-Ig



Basic Information

Catalog Number:

68717-1-Ig

Size:

150ul, Concentration: 1000 ug/ml by Nanodrop;

Source:

Mouse

Isotype:

IgG2a

Immunogen Catalog Number:

AG15263

GenBank Accession Number:

BC031672

GeneID (NCBI):

79932

UNIPROT ID:

Q8IZAO

Full Name:

KIAA0319-like

Calculated MW:

1049 aa, 116 kDa

Observed MW:

140-150 kDa

Purification Method:

Protein A purification

CloneNo.:

2E11D9

Recommended Dilutions:

WB 1:5000-1:50000

Applications

Tested Applications:

WB, ELISA

Species Specificity:

Human

Positive Controls:

WB : A549 cells, LNCaP cells, HeLa cells, HEK-293 cells, HepG2 cells, K-562 cells, U2OS cells

Background Information

KIAA0319L (also known as AAVR) is a type-I transmembrane protein that acts as a receptor for adeno-associated virus (AAV) and is involved in adeno-associated virus infection through endocytosis system (PMID: 26814968). It has been shown that KIAA0319L is highly-glycosylated and can localise to the plasma membrane, endosomes, Golgi and trans-Golgi network (TGN) (PMID: 29991750). KIAA0319L gene has been associated with susceptibility to developmental dyslexia or reading disabilities (PMID: 19085271). The predicted molecular mass of KIAA0319L is 116 kDa, but western blot detected protein at 150 kDa in lysate from HeLa cells, likely due to glycosidic linkages (PMID: 28679762; 30962536).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68717-1-Ig (KIAA0319L antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.