For Research Use Only

uPAR/CD87 Monoclonal antibody

Catalog Number: 68727-1-Ig



Basic Information

Catalog Number: GenBank Accession Number:

68727-1-lg BC002788 GeneID (NCBI):

Nanodrop: **UNIPROT ID:** Q03405 Mouse Full Name: Isotype: **PLAUR**

150ul, Concentration: 1000 ug/ml by 5329

lgG2b Calculated MW: Immunogen Catalog Number: 37 kDa AG28056 Observed MW:

35-50 kDa

Purification Method:

Protein A purification CloneNo.:

2A3E8

Recommended Dilutions: WB 1:1000-1:4000 IHC 1:250-1:1000

Applications

Tested Applications: WB, IHC, ELISA Species Specificity:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: U2OS cells, U-251 cells, A549 cells IHC: human lung squamous cell cancer,

Background Information

uPAR is a highly glycosylated, GPI-anchored membrane protein. In addition to the membrane-anchored form, uPAR is released from the plasma membrane by cleavage of the GPI anchor and can be found as a soluble form (suPAR). uPAR contains three homologous domains (D1-D3) of which the N-terminal one (D1) represents the uPA-binding domain. After binding to uPAR, uPA cleaves plasminogen, generating the active protease plasmin which is involved in a wide variety of physiologic and pathologic processes. In addition to regulating proteolysis, uPAR has important function in cell adhesion, migration and proliferation. Studies reveal that uPAR expression is elevated during inflammation and tissue remodeling and in many human cancers, in which it frequently indicates poor prognosis. (PMID 20027185; 12461559)

Storage

Storage:


Store at -20°C. Stable for one year after shipment. Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



U2OS cells were subjected to SDS PAGE followed by western blot with 68727-1-1g (uPAR/CD87 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human lung squamous cell cancer slide using 68727-1-1g (uPAR/CD87 antibody) at dilution of 1:500 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).