

For Research Use Only

Mono/Di-Methyl-Histone H3 (Lys9) Monoclonal antibody

Catalog Number: 68825-1-Ig



Basic Information

Catalog Number:

68825-1-Ig

Size:

150ul, Concentration: 1000 ug/ml by
Nanodrop;

Source:

Mouse

Isotype:

IgG2a

GenBank Accession Number:

BC066245

GeneID (NCBI):

8350

UNIPROT ID:

P68431

Full Name:

histone cluster 1, H3a

Observed MW:

17 kDa

Purification Method:

Protein A purification

CloneNo.:

2E6D11

Recommended Dilutions:

WB 1:5000-1:50000

Applications

Tested Applications:

WB, ELISA

Species Specificity:

Human, mouse, rat

Positive Controls:

WB : HeLa cells, HEK-293 cells, LNCaP cells, Jurkat cells, MOLT-4 cells, K-562 cells, NIH/3T3 cells, C2C12 cells, HSC-T6 cells

Background Information

Histones, including H1/H5 (linker histones), H2, H3, and H4 (core histones), are nucleic proteins which interact with DNA to form the nucleosomes and play important roles in gene regulation and DNA replication. Histone proteins are highly post-translationally modified while Histone H3 is the most extensively modified. Methylation of Histone H3 at lysine 9 is linked to transcriptional repression. This antibody is specific to monomethyl or dimethyl-Histone H3 while it does not recognize trimethyl-Histone H3 (Lys9). It is also named as H3K9me1/2.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68825-1-Ig (Histone H3.1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.