## For Research Use Only

## PARK7/DJ-1 Monoclonal antibody

Catalog Number: 68915-1-Ig



**Basic Information** 

Catalog Number: GenBank Accession Number:

68915-1-lg BC008188
Size: Genel D (NCBI):

150ul , Concentration: 500 ug/ml by 11315
Nanodrop; UNIPROT ID:
Source: Q99497
Mouse Full Name:

Isotype: Parkinson disease (autosomal IgG1 recessive, early onset) 7

Immunogen Catalog Number:Calculated MW:AG28526189 aa, 20 kDaObserved MW:

20-25 kDa

**Applications** 

**Tested Applications:** 

WB, ELISA

Species Specificity: human, pig, rabbit

Positive Controls:

WB: HeLa cells, pig brain tissue, HEK-293 cells, Jurkat

**Purification Method:** 

CloneNo.:

4E10B2

Protein G purification

Recommended Dilutions:

WB 1:2000-1:10000

cells, rabbit brain tissue

## **Background Information**

PARK7, also named as DJ1, belongs to the peptidase C56 family. It protects cells against oxidative stress and cell death. PARK7 plays a role in regulating expression or stability of the mitochondrial uncoupling proteins SLC25A14 and SLC25A27 in dopaminergic neurons of the substantia nigra pars compacta and attenuates the oxidative stress induced by calcium entry into the neurons via L-type channels during pacemaking. It eliminates hydrogen peroxide and protects cells against hydrogen peroxide-induced cell death. PARK7 has cell-growth promoting activity and transforming activity. It may function as a redox-sensitive chaperone. It's precursor undergoes a cleavage of a C-terminal peptide and subsequent activation of protease activity in response to oxidative stress. The amino acid replace at 166 (L — P) reduces PARK7 protein stability and leads to increased degradation. The predicted MW of this protein is 20 kDa, An additional 25 kDa band can be observed due to modification (PMID: 31767755).

Storage

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68915-1-1g (PARK7/DJ-1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 68915-1-1g (PARK7/DJ-1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.