For Research Use Only

NeutraKine® IL-3 Mouse McAb

Catalog Number:69004-1-Ig



Basic Information	Catalog Number: 69004-1-lg Size: 100ug Source: Mouse Isotype: IgG1 Immunogen Catalog Number: HZ-1074	GenBank Accession Number: GeneID (NCBI): 3562 Full Name: interleukin 3 (colony-stimulating factor, multiple)	Purification Method: Protein G purification CloneNo.: 3C11G7
Applications	Tested Applications: Neutralization, ELISA Species Specificity: human	Positive Con Neutralizatio	
Background Information	Interleukin-3 (IL-3) is a multilineage hematopoietic growth factor that promotes the proliferation, differentiation and survival of early multilineage hematopoietic progenitors. In particular, this cytokine plays a key role in stimulating the proliferation and survival of myeloid precursors. It is involved in a variety of cell activities such as cell growth, differentiation and apoptosis. This cytokine has been shown to also possess neurotrophic activity, and it may be associated with neurologic disorders (PMID: 24103757;2698876;2809196). This antibody can be used to neutralize the bioactivity of IL-3.		
Storage	Storage: Lyophilized antibodies are stable for 1 year from the date of receipt if stored between (-20°C) and (-80°C). Upon reconstitution we recommend that the solution can be stored at (4°C) for short term or at (-20°C) to (-80°C) for long term. Repeated freeze thaw cycles should be avoided with reconstituted products. Storage Buffer: Sterile PBS, pH7.4 Aliquoting is unnecessary for -20°C storage		
*** 20ul sizes contain 0.1% BSA		-	

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Recombinant human IL-3 (Cat.NO. HZ-1074) stimulates proliferation of TF-1 cell line (human erythroleukemic cell line) in a dose-dependent manner (blue curve, refer to bottom X-left Y). The activity of human IL-3 (10 ng/mL HZ-1074) is neutralized by mouse anti-human IL-3 monoclonal antibody 69004-1-1g at serial dose (red curve, refer to top X-right Y). The ND50 is typically 40-70 ng/mL

Recombinant human IL-3 (Cat.NO. HZ-1074) stimulates proliferation of TF-1 cell line (human erythroleukemic cell line) in a dose-dependent manner (blue curve, refer to bottom X-left Y). The activity of human IL-3 (0.5 ng/mL HZ-1074) is neutralized by mouse anti-human IL-3 monoclonal antibody 69004-1-lg at serial dose (red curve, refer to top X-right Y). The ND50 is typically 0.8-2.4 ng/mL