For Research Use Only

NeutraKine® IL-3 Monoclonal antibody, proteintech® PBS Only


Catalog Number:69004-1-PBS

Basic Information	Catalog Number: 69004-1-PBS Size: 100ug , Concentration: 1mg/ml by Nanodrop; Source: Mouse Isotype: IgG1 Immunogen Catalog Number: HZ-1074	GenBank Accession Number: GeneID (NCBI): 3562 Full Name: interleukin 3 (colony-stimulating factor, multiple)	Purification Method: Protein G purification CloneNo.: 3C11G7
Applications	Tested Applications: Neutralization, ELISA Species Specificity: human		
Background Information	Interleukin-3 (IL-3) is a multilineage hematopoietic growth factor that promotes the proliferation, differentiation and survival of early multilineage hematopoietic progenitors. In particular, this cytokine plays a key role in stimulating the proliferation and survival of myeloid precursors. It is involved in a variety of cell activities such as cell growth, differentiation and apoptosis. This cytokine has been shown to also possess neurotrophic activity, and it may be associated with neurologic disorders (PMID: 24103757;2698876;2809196). This antibody can be used to neutralize the bioactivity of IL-3.		
Storage	This antibody can be used to neutral Storage: Store at -80°C. Storage Buffer: PBS only, pH7.3	lize the bloactivity of IL-3.	


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Recombinant human IL-3 (Cat.NO. HZ-1074) stimulates proliferation of TF-1 cell line (human erythroleukemic cell line) in a dosedependent manner (blue curve, refer to bottom X-left Y). The activity of human IL-3 (10 ng/mL HZ-1074) is neutralized by mouse anti-human IL-3 monoclonal antibody 69004-1-Ig at serial dose (red curve, refer to top X-right Y). The ND50 is typically 40-70 ng/mL. This data was developed using the same antibody clone with 69004-1-PBS in a



Recombinant human IL-3 (Cat.NO. HZ-1074) stimulates proliferation of TF-1 cell line (human erythroleukemic cell line) in a dosedependent manner (blue curve, refer to bottom X-left Y). The activity of human IL-3 (0.5 ng/mL HZ-1074) is neutralized by mouse anti-human IL-3 monoclonal antibody 69004-1-Ig at serial dose (red curve, refer to top X-right Y). The ND50 is typically 0.8-2.4 ng/mL This data was developed using the same antibody clone with 69004-1-PBS in a