## For Research Use Only

## NeutraKine® IFN Alpha 2A Monoclonal antibody

Catalog Number:69008-1-lg




Basic Information	Catalog Number: 69008-1-lg Size: 100ug Source: Mouse	GenBank Accession Number:	Purification Method: Protein G purification CloneNo.: 2H8G1			
		GeneID (NCBI): 3440 Full Name: interferon, alpha 2				
				lsotype:		
				lgG1		
				Immunogen Catalog Number: HZ-1066		
				Applications	Tested Applications:	
	ELISA, Neutralization					
	Species Specificity: Human					
	Packground Information	Interferon alpha 2 is a type 1 interf	feron Interferon alpha 2 is secreted b	y virus-infected cells and inhibits other cells		
Background Information	from further infection. Recombinant human interferon alpha 2 has been approved for therapeutic application in a range of human oncological and viral diseases. Three Interferon alpha 2 subvariants (2A, 2B, 2C) differ by only one or two amino acids at positions 23 and/or 34 of the mature protein (PMID: 25982860; 10672347; 1694761).					
	This antibody can be used to neutralize the bioactivity of Interferon alpha 2.					
Storage	Storage:					
	reconstitution we recommend that	, ,	ored between (-20°C) and (-80°C). Upon r short term or at (-20°C) to (-80°C) for long I products.			
	Aliquoting is unnecessary for -20 $^{\circ}$	C storage				
*** 20ul sizes contain 0.1% BSA						

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free<br/>in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Recombinant human IFN alpha 2A (Cat.NO. HZ-1066) inhibits the growth of TF-1 cell line (human erythroleukemic cell line) in a dose-dependent manner (blue curve, refer to bottom X-left Y). The activity of human IFN alpha 2A (100 ng/mL HZ-1066) is neutralized by mouse anti-human IFN alpha 2A monoclonal antibody 69008-1-Ig at serial dose (red curve, refer to top X-right Y). The ND50 is typically 300-500 ng/mL