For Research Use Only

NeutraKine®FGF basic Mouse McAb

Catalog Number:69024-1-lg 1 Publications



Basic Information	Catalog Number:	GenBank Accession Num	nber:	Purification Method:	
	69024-1-lg	GeneID (NCBI):		Protein G purification	
	Size:	2247		CloneNo.:	
	100ug	Full Name:	- (1 - 1)	3C9C3	
	Source: Mouse	fibroblast growth factor 2 (basic)			
	lsotype: lgG1				
	Immunogen Catalog Number: HZ-1285				
Applications	Tested Applications:	P	Positive Controls: Neutralization : BALB/3T3 cells,		
	ELISA, Neutralization Cited Applications: Cell treatment	Ν			
	Species Specificity: Human				
	Cited Species: human				
Background Information), also referred to as basic FGI	F, belongs to	a family of growth factors that stimul	
Background Informatior	Fibroblast growth factor-2 (FGF-2 proliferation of cells of mesenchy multiple polyadenylation sites, a resulting in at least five different	mal, epithelial and neuroect nd is alternatively translated isoforms with distinct proper	odermal orig I from non-Al rties.	a family of growth factors that stimul in. The mRNA for this gene contains JG (CUG) and AUG initiation codons,	
Background Informatior	Fibroblast growth factor-2 (FGF-2 proliferation of cells of mesenchy multiple polyadenylation sites, a	mal, epithelial and neuroect nd is alternatively translated isoforms with distinct proper	odermal orig I from non-Al rties.	in. The mRNA for this gene contains	
	Fibroblast growth factor-2 (FGF-2 proliferation of cells of mesenchy multiple polyadenylation sites, a resulting in at least five different This antibody is used to neutraliz	mal, epithelial and neuroect nd is alternatively translated isoforms with distinct proper	odermal orig I from non-Al rties.		
	Fibroblast growth factor-2 (FGF-2 proliferation of cells of mesenchy multiple polyadenylation sites, a resulting in at least five different This antibody is used to neutraliz	mal, epithelial and neuroect nd is alternatively translated isoforms with distinct proper e the bioactivity of FGFbasic- Pubmed ID Journal	odermal orig I from non-Al rties.	rin. The mRNA for this gene contains JG (CUG) and AUG initiation codons, Application	
Background Information Notable Publications Storage	Fibroblast growth factor-2 (FGF-2 proliferation of cells of mesenchy multiple polyadenylation sites, a resulting in at least five different This antibody is used to neutraliz Author F Dandan Li Storage: Lyophilized antibodies are stable	mal, epithelial and neuroect nd is alternatively translated isoforms with distinct proper e the bioactivity of FGFbasic- Pubmed ID Journal 38479160 Int Immur for 1 year from the date of re at the solution can be stored a	odermal orig d from non-Al rties. -TS. nopharmacol eccipt if store tt(4°C) for sho	rin. The mRNA for this gene contains JG (CUG) and AUG initiation codons, Application Cell treatment d between (-20°C) and (-80°C). Upon ort term or at(-20°C) to (-80°C) for long	
Notable Publications	Fibroblast growth factor-2 (FGF-2 proliferation of cells of mesenchy multiple polyadenylation sites, a resulting in at least five different This antibody is used to neutraliz Author P Dandan Li 2 Storage: Lyophilized antibodies are stable reconstitution we recommend tha term. Repeated freeze thaw cycle Storage Buffer:	mal, epithelial and neuroect nd is alternatively translated isoforms with distinct proper e the bioactivity of FGFbasic- Pubmed ID Journal 58479160 Int Immur for 1 year from the date of re it the solution can be stored a s should be avoided with reco	odermal orig d from non-Al rties. -TS. nopharmacol eccipt if store tt(4°C) for sho	rin. The mRNA for this gene contains JG (CUG) and AUG initiation codons, Application Cell treatment d between (-20°C) and (-80°C). Upon ort term or at(-20°C) to (-80°C) for long	
Notable Publications	Fibroblast growth factor-2 (FGF-2 proliferation of cells of mesenchy multiple polyadenylation sites, a resulting in at least five different This antibody is used to neutraliz Author P Dandan Li 2 Storage: Lyophilized antibodies are stable reconstitution we recommend that term. Repeated freeze thaw cycle Storage Buffer: Sterile PBS, pH7.4	mal, epithelial and neuroect nd is alternatively translated isoforms with distinct proper e the bioactivity of FGFbasic- Pubmed ID Journal 58479160 Int Immur for 1 year from the date of re it the solution can be stored a s should be avoided with reco	odermal orig d from non-Al rties. -TS. nopharmacol eccipt if store tt(4°C) for sho	rin. The mRNA for this gene contains JG (CUG) and AUG initiation codons, Application Cell treatment d between (-20°C) and (-80°C). Upon ort term or at(-20°C) to (-80°C) for long	

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Recombinant human FGF Basic (Cat.NO. HZ-1285) stimulates the proliferation of BALB/3T3 cells (mouse fibroblast cell) at low FBS condition in a dose-dependent manner (blue curve, refer to bottom X-left Y axis). The activity of human FGF Basic (0.1 ng/mL HZ-1285) is neutralized by mouse anti-human FGF Basic monoclonal antibody 69024-1-lg at serial dose (red curve, refer to top X-right Y axis). The ND50 is typically 1-10 µg/mL. Mouse IgG1 Isotype Control monoclonal