## For Research Use Only

## NeutraControl IL-6 Mouse McAb

Catalog Number:69501-1-lg 1 Publications


Antibodies | ELISA kits | Proteins www.ptglab.com


Basic Information	Catalog Number:	GenBank Accession Number:	Purification Method:	
	69501-1-lg	GeneID (NCBI):	Protein G purification	
	Size:	3569	CloneNo.:	
	100ug	ENSEMBL Gene ID:	1E9B8	
	Source:	ENSG0000136244	Recommended Dilutions:	
	Mouse	Full Name:	Non-Neutralization: 1:10-1:100 ELISA: 1:10-1:100	
	Isotype: IgG1	interleukin 6 (interferon, beta 2)	ELISA. 1.10-1.100	
	Immunogen Catalog Number: HZ-1019			
Applications	Tested Applications:	Positive Controls:		
TT	ELISA, Non-Neutralization	Non-Neut	tralization : hybridoma,	
	Cited Applications: ELISA	ELISA : Recombinant protein,		
	Species Specificity: human			
	Cited Species:			
Background Information	human Interleukin-6 (IL-6) is an interleuk	(in that acts as both a pro-inflammate	ry and anti-inflammatory cytokine. IL-6	
Background Information	Interleukin-6 (IL-6) is an interleuk protein is secreted by a variety of glycosylated molecule. IL-6 plays involved in lymphocyte and mom nerve cells differentiation Acts or IL-6 is also considered a myokine contraction. IL-6 has been shown	cell types including T cells and macr s an essential role in the final differer ocyte differentiation. It induces myel n B-cells, T-cells, hepatocytes, hemato , a cytokine produced from muscle, ar to interact with interleukin-6 receptor metabolism, and cancer progression,	ophages as phosphorylated and variably itiation of B-cells into Ig-secreting cells oma and plasmacytoma growth and induce poietic progenitor cells and cells of the CN nd is elevated in response to muscle	
Background Information	Interleukin-6 (IL-6) is an interleuk protein is secreted by a variety of glycosylated molecule. IL-6 plays involved in lymphocyte and mon- nerve cells differentiation Acts or IL-6 is also considered a myokine contraction. IL-6 has been shown t involved in hematopoiesis, bone directing transition from innate to	cell types including T cells and macr an essential role in the final differen ocyte differentiation. It induces myel n B-cells, T-cells, hepatocytes, hemato , a cytokine produced from muscle, ar to interact with interleukin-6 receptor metabolism, and cancer progression, o acquired immunity.	ophages as phosphorylated and variably itiation of B-cells into Ig-secreting cells oma and plasmacytoma growth and induce poietic progenitor cells and cells of the CN nd is elevated in response to muscle and glycoprotein 130. Additionally, IL-6 is	
Background Information	Interleukin-6 (IL-6) is an interleuk protein is secreted by a variety of glycosylated molecule. IL-6 plays involved in lymphocyte and mon- nerve cells differentiation Acts or IL-6 is also considered a myokine contraction. IL-6 has been shown t involved in hematopoiesis, bone directing transition from innate to This antibody is a neutralizing co could not neutralize human IL-6.	cell types including T cells and macr an essential role in the final differen ocyte differentiation. It induces myel n B-cells, T-cells, hepatocytes, hemato , a cytokine produced from muscle, ar to interact with interleukin-6 receptor metabolism, and cancer progression, o acquired immunity.	ophages as phosphorylated and variably itiation of B-cells into Ig-secreting cells oma and plasmacytoma growth and induce poietic progenitor cells and cells of the CN nd is elevated in response to muscle and glycoprotein 130. Additionally, IL-6 is and has been defined an essential role in	
	<ul> <li>Interleukin-6 (IL-6) is an interleuk protein is secreted by a variety of glycosylated molecule. IL-6 plays involved in lymphocyte and monnerve cells differentiation Acts or IL-6 is also considered a myokine contraction. IL-6 has been shown to involved in hematopoiesis, bone directing transition from innate to This antibody is a neutralizing concould not neutralize human IL-6.</li> <li>Author</li> </ul>	cell types including T cells and macr an essential role in the final differer ocyte differentiation. It induces myel B-cells, T-cells, hepatocytes, hemato , a cytokine produced from muscle, ar to interact with interleukin-6 receptor metabolism, and cancer progression, o acquired immunity. ntrol antibody for IL-6, the immunoge	ophages as phosphorylated and variably itiation of B-cells into Ig-secreting cells oma and plasmacytoma growth and induce poietic progenitor cells and cells of the CN id is elevated in response to muscle and glycoprotein 130. Additionally, IL-6 is and has been defined an essential role in n is the same as Neutrakine 69001-1-lg bur Application	
	<ul> <li>Interleukin-6 (IL-6) is an interleuk protein is secreted by a variety of glycosylated molecule. IL-6 plays involved in lymphocyte and more nerve cells differentiation Acts or IL-6 is also considered a myokine contraction. IL-6 has been shown to involved in hematopoiesis, bone directing transition from innate to This antibody is a neutralizing concould not neutralize human IL-6.</li> <li>Author</li> <li>Yao Zhang</li> <li>Storage:</li> <li>Lyophilized antibodies are stable reconstitution we recommend that the stable reconstitution we recommend the stable reconstitution we recommend that the stable recon</li></ul>	icell types including T cells and macristan essential role in the final differency ocyte differentiation. It induces myells a B-cells, T-cells, hepatocytes, hemator, a cytokine produced from muscle, are to interact with interleukin-6 receptor metabolism, and cancer progression, o acquired immunity.         ntrol antibody for IL-6, the immunoge         Pubmed ID       Journal         36356714       Free Radic Biol N         for 1 year from the date of receipt if s the solution can be stored at(4°C) for s should be avoided with reconstitute	ophages as phosphorylated and variably titation of B-cells into Ig-secreting cells oma and plasmacytoma growth and induce poietic progenitor cells and cells of the CN id is elevated in response to muscle and glycoprotein 130. Additionally, IL-6 is and has been defined an essential role in in is the same as Neutrakine 69001-1-1g bu Application field ELISA	

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data





Recombinant human IL-6 (Cat.NO. HZ-1019) stimulates proliferation of hybridoma (Proteintech mouse anti-GST clone 3G12B10) in a dose-dependent manner (blue curve, refer to bottom X-left Y axis). The activity of human IL-6 (1 ng/mL HZ-1019) is neutralized by mouse anti-human IL-6 monoclonal antibody 69001-1-1g at serial dose (red curve, refer to top X-right Y axis). The ND50 is typically 10-30 ng/mL The NeutraControl mouse anti-human IL-6 monoclonal antibody Indirect ELISA was carried out by coating recombinant Human IL-6 (Cat.NO. HZ-1019) at 70 ng/well followed by blocking and adding serial diluted IL-6 antibody 69001-1-Ig and 69501-1-Ig respectively. Signal was developed with TMB and stopped by H2SO4. Signal strength was measured by absorbance at 450 nm.