

For Research Use Only

NeutraControl IL-17A Monoclonal antibody



Catalog Number: 69521-1-Ig

Basic Information

Catalog Number:

69521-1-Ig

Size:

100ug

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

HZ-1113

GenBank Accession Number:

GeneID (NCBI):

3605

ENSEMBL Gene ID:

ENSG00000112115

Full Name:

interleukin 17A

Purification Method:

Protein G purification

CloneNo.:

2A7E2

Applications

Tested Applications:

ELISA, Non-Neutralization

Species Specificity:

Human

Background Information

IL17A, also named as IL-17, is a proinflammatory cytokine. IL-17, synthesized only by memory T cells and natural killer cells, has pleiotropic effects, mainly in the recruitment and activation of neutrophils. This cytokine regulates the activities of NF-kappaB and mitogen-activated protein kinases. This cytokine can stimulate the expression of IL6 and cyclooxygenase-2 (PTGS2/COX-2), as well as enhance the production of nitric oxide (NO). High levels of this cytokine are associated with several chronic inflammatory diseases including rheumatoid arthritis, psoriasis and multiple sclerosis. The IL-17 receptor is a type I transmembrane protein, that is widely expressed on epithelial cells, fibroblasts, B and T cells, and monocytic cells. In psoriatic skin lesions, both Th17 cells and their downstream effector molecules, e.g. IL-17 and IL-22, are highly increased.

This antibody is a neutralizing control antibody for IL-17A, the immunogen is the same as Neutrakine 69021-1-Ig but could not neutralize human IL-17A.

Storage

Storage:

Lyophilized antibodies are stable for 1 year from the date of receipt if stored between (-20°C) and (-80°C). Upon reconstitution we recommend that the solution can be stored at(4°C) for short term or at(-20°C) to (-80°C) for long term. Repeated freeze thaw cycles should be avoided with reconstituted products.

Storage Buffer:

Sterile PBS.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA



For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Recombinant human IL-17A (Cat.NO. HZ-1113) stimulates HT-1080 cells (human fibroblast cell line) produce IL-6 in a dose dependent manner (blue curve, refer to bottom X-Left Y axis). The activity of human IL-17A (40ng/mL HZ-1113) is neutralized by mouse anti-human IL-17A monoclonal antibody 69021-1-Ig (red curve, refer to top X-right Y axis). The ND50 is typically 200-600ng/mL. The NeutraControl mouse anti-human IL-17A monoclonal antibody 69521-1-Ig could recognize human

Indirect ELISA was carried out by coating recombinant Human IL-17A (Cat.NO. HZ-1113) at 70 ng/well followed by blocking and adding serial diluted IL-17A antibody 69021-1-Ig and 69521-1-Ig respectively. Signal was developed with TMB and stopped by H2SO4. Signal strength was measured by absorbance at 450 nm.