## For Research Use Only

## NeutraControl HGF Monoclonal antibody



Catalog Number: 69527-1-Ig

**Basic Information** 

Catalog Number:

69527-1-lg

Size: 100ug

Source: Mouse

Isotype: IgG2a

Immunogen Catalog Number:

HZ-1084

GenBank Accession Number:

GeneID (NCBI):

3082

hepatocyte growth factor (hepapoietin A; scatter factor)

Full Name:

**Purification Method:** 

Protein A purification CloneNo.:

2G4G8

**Applications** 

**Tested Applications:** 

ELISA, Non-Neutralization

Species Specificity:

Human

## **Background Information**

Hepatocyte growth factor (HGF) is the most potent mitogen of mature hepatocytes in primary culture. HGF is derived from a biologically inactive single chain precursor of 728 amino acids (pro-HGF) localized mostly on the cell surface and in the extracellular matrix. The mature form produced following proteolytic cleavage is composed of a 69-kDa a-subunit (containing four kringle domains) and the 34 kDa β-subunit, similar to the catalytic domain of serine proteases, but with amino acid substitutions in the active site. HGF is a pleiotropic cytokine which exerts a variety of effects on several cells, being involved in the regulation of many biological processes, such as inflammation, tissue repair, morphogenesis, angiogenesis, tumour propagation, immunomodulation of viral infections and cardiometabolic activities.

This antibody is a neutralizing control antibody for HGF, the immunogen is the same as Neutrakine 69027-1-Ig but could not neutralize human HGF.

Storage

Lyophilized antibodies are stable for 1 year from the date of receipt if stored between (-20°C) and (-80°C). Upon reconstitution we recommend that the solution can be stored at (4°C) for short term or at (-20°C) to (-80°C) for long term. Repeated freeze thaw cycles should be avoided with reconstituted products.


Storage Buffer:


Sterile PBS.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

## **Selected Validation Data**





Recombinant human HGF (Cat.NO. HZ-1084) stimulates Saos-2 cells (human osteosarcoma cell) produce IL-11 in a dose dependent manner (blue curve, refer to bottom X-Left Y axis). The activity of human HGF (50ng/mL HZ-1084) is neutralized by mouse anti-human HGF monoclonal antibody 69027-1-1g (red curve, refer to top X-right Y axis). The ND50 is typically 2-8µg/mLThe NeutraControl mouse anti-human HGF antibody 69527-1-1g could recognize human HGF but could not neutralize

Indirect ELISA was carried out by coating recombinant Human HGF (Cat.NO. HZ-1084) at 70 ng/well followed by blocking and adding serial diluted Thrombopoietin antibody 69027-1-lg and 69527-1-lg respectively. Signal was developed with TMB and stopped by H2SO4. Signal strength was measured by absorbance at 450 nm.