NeutraControl IL-12 p35 Monoclonal proteintech antibody


Catalog Number:69529-1-lg

Basic Information	Catalog Number: 69529-1-lg Size: 100ug Source: Mouse Isotype: IgG1 Immunogen Catalog Number: HZ-1256	GenBank Accession Number: GeneID (NCBI): 3592 Full Name: interleukin 12A (natural killer cell stimulatory factor 1, cytotoxic lymphocyte maturation factor 1, p35	Purification Method: Protein G purification CloneNo.: 2C4G2
Applications	Tested Applications: ELISA, Non-Neutralization Species Specificity: Human		
Background Information	IL12 p35 is a subunit of a cytokine that acts on T and natural killer cells, and has a broad array of biological activities. The cytokine is a disulfide-linked heterodimer composed of the 35-kD subunit encoded by this gene, and a 40-kD subunit that is a member of the cytokine receptor family. This cytokine is required for the T-cell-independent induction of IFN-gamma, and is important for the differentiation of both Th1 and Th2 cells. The responses of lymphocytes to this cytokine are mediated by the activator of transcription protein STAT4. Nitric oxide synthase 2A (NOS2A/NOS2) is found to be required for the signaling process of this cytokine in innate immunity. This antibody is a neutralizing control antibody for IL-12 p35, the immunogen is the same as Neutrakine 69029-1-lg but could not neutralize human IL-12 p35.		
Storage *** 20ul sizes contain 0.1% BSA	reconstitution we recommend that the	r 1 year from the date of receipt if store he solution can be stored at(4°C) for sho hould be avoided with reconstituted pr storage	ort term or at(-20°C) to (-80°C) for long


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Recombinant human IL-12 (Cat.NO. HZ-1256) stimulates NK-92 cells produce IFN gamma in a dose dependent manner (blue curve, refer to bottom X-Left Y axis). The activity of human IL-12 (10 ng/mL HZ-1256) is neutralized by mouse anti-human IL-12 p35 monoclonal antibody 69029-1-1g (red curve, refer to top Xright Y axis). The ND50 is typically 0.5-1 µg/mL. The control mouse anti-human IL-12 p35 monoclonal antibody 69529-1-1g could recognize human IL-12, but could not



Indirect ELISA was carried out by coating recombinant Human IL-12 (Cat.NO. HZ-1256) and IL-12 p40 (Cat.NO. HZ-1321) respectively at 70 ng/well followed by blocking and adding serial diluted 69029-1-1g, 69529-1-1g and 69006-1-1g respectively. HRP-goat antimouse was used for detection. Signal was developed with TMB and stopped by H2SO4. Signal strength was measured by absorbance at 450 nm. The result suggests that 69029-1-1g and 69529-1-1g only recognize IL-12(p35 and