

NeutraControl FLT3 Ligand Monoclonal antibody

Catalog Number: 69532-1-Ig

Basic Information

Catalog Number:

69532-1-Ig

Size:

100ug

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

HZ-1151

GenBank Accession Number:

GeneID (NCBI):

2323

Full Name:

fms-related tyrosine kinase 3 ligand

Purification Method:

Protein G purification

CloneNo.:

3C9C7

Applications

Tested Applications:

ELISA, Non-Neutralization

Species Specificity:

Human

Background Information

The FMS-related tyrosine kinase 3 (FLT3) ligand (FLT3LG), a growth factor, is mainly derived from lymphocytes in the tumor microenvironment. It binds to FLT3 on DCs to enhance their differentiation and expansion. Therefore, it may assist DCs in tumor antigen presentation and antitumor immune response. In addition, mature DCs express FLT3 and proliferate massively under the trigger of FLT3 Ligand.

This antibody is a neutralizing control antibody for FLT3 Ligand, the immunogen is the same as Neutrakine 69032-Ig but could not neutralize human FLT3 Ligand.

Storage

Storage:

Lyophilized antibodies are stable for 1 year from the date of receipt if stored between (-20°C) and (-80°C). Upon reconstitution we recommend that the solution can be stored at 4°C for short term or at (-20°C) to (-80°C) for long term. Repeated freeze thaw cycles should be avoided with reconstituted products.

Storage Buffer:

Sterile PBS.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA



For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Recombinant human FLT3 Ligand (Cat.NO. HZ-1151) stimulates proliferation of OCI-AML5 cells in a dose-dependent manner (blue curve, refer to bottom X-left Y). The activity of human FLT3 (1 ng/mL) is neutralized by mouse anti-human FLT3 Ligand monoclonal antibody 69032-1-Ig at serial dose (red curve, refer to top X-right Y). The ND50 is typically 0.01-0.05 µg/mL. The control mouse anti-human FLT3 Ligand monoclonal antibody 69532-1-Ig could not recognize human FLT3 Ligand, but could not

Indirect ELISA was carried out by coating recombinant Human FLT3 Ligand (Cat.NO. HZ-1151) at 70 ng/well followed by blocking and adding serial diluted FLT3 Ligand antibody 69032-1-Ig and 69532-1-Ig respectively. Signal was developed with TMB and stopped by H2SO4. Signal strength was measured by absorbance at 450 nm.