NeutraControl M-CSF Monoclonal antibody

Catalog Number:69533-1-lg





Basic Information	Catalog Number: 69533-1-lg Size: 100ug Source: Mouse Isotype: IgG1 Immunogen Catalog Number: HZ-1192	GenBank Accession Number: GeneID (NCBI): 1435 Full Name: colony stimulating factor 1 (macrophage)	Purification Method: Protein G purification CloneNo.: 4B3C1
Applications	Tested Applications: ELISA, Non-Neutralization Species Specificity: Human		
Background Information	M-CSF plays an essential role in the regulation of survival, proliferation, and differentiation of hematopoietic precursor cells, especially mononuclear phagocytes. M-CSF promotes the release of proinflammatory chemokines and thereby plays an important role in innate immunity and in inflammatory processes. It also plays an important role in the regulation of osteoclast proliferation and differentiation, the regulation of bone resorption, and is required for normal bone development. On the cellular level, M-CSF promotes reorganization of the actin cytoskeleton, regulates the formation of membrane ruffles, cell adhesion, and cell migration, and plays a role in lipoprotein clearance.		
Storage *** 20ul sizes contain 0.1% BSA	reconstitution we recommend that	or 1 year from the date of receipt if st the solution can be stored at(4°C) for should be avoided with reconstituted	ored between (-20°C) and (-80°C). Upon short term or at(-20°C) to (-80°C) for long I products.

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Recombinant human M-CSF(Cat.NO. HZ-1192) stimulates proliferation of OCI-AML5 cells in a dose-dependent manner (blue curve, refer to bottom X-left Y). The activity of human M-CSF (5 ng/mL) is neutralized by mouse anti-human M-CSF monoclonal antibody 69033-1-Ig at serial dose (red curve, refer to top X-right Y). The ND50 is typically 0.1-0.4 µg/mL. The control mouse anti-human M-CSF monoclonal antibody 69533-1-Ig could recognize human M-CSF, but could not neutralize human M-CSF

Indirect ELISA was carried out by coating recombinant Human M-CSF (Cat.NO. HZ-1192) at 70 ng/well followed by blocking and adding serial diluted M-CSF antibody 69033-1-1g and 69533-1-1g respectively. Signal was developed with TMB and stopped by H2SO4. Signal strength was measured by absorbance at 450 nm.