

For Research Use Only

Phospho-Beta Catenin (Ser552) Recombinant antibody, PBS Only

Catalog Number: 80035-2-PBS



Basic Information

Catalog Number:

80035-2-PBS

Size:

100ug, Concentration: 1 mg/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

BC058926

GeneID (NCBI):

1499

ENSEMBL Gene ID:

ENSG00000168036

UNIPROT ID:

P35222

Full Name:

catenin (cadherin-associated protein), beta 1, 88kDa

Calculated MW:

781 aa, 86 kDa

Observed MW:

86-100 kDa

Purification Method:

Protein A purification

CloneNo.:

250272E1

Applications

Tested Applications:

WB, Indirect ELISA

Species Specificity:

human

Background Information

β -Catenin, also known as CTNNB1, is an evolutionarily conserved, multifunctional intracellular protein. β -Catenin is a 92-kDa protein, originally identified in cell adherens junctions (AJs) where it functions to bridge the cytoplasmic domain of cadherins to α -catenin and the actin cytoskeleton. Besides its essential role in the AJs, β -catenin is also a key downstream component of the canonical Wnt pathway that plays diverse and critical roles in embryonic development and adult tissue homeostasis. The Wnt/ β -catenin pathway is also involved in the activation of other intracellular messengers such as calcium fluxes, JNK, and SRC kinases. Deregulation of β -catenin activity is associated with multiple diseases including cancers (PMID: 22617422, PMID: 18334222). Both AMPK and AKT phosphorylates β -catenin at Ser 552. The phosphorylation of β -catenin at Ser552 can stabilize β -catenin, enhance its nuclear accumulation and increase its transcriptional activity (PMID: 2865606, PMID: 20361929).

Storage

Storage:

Store at -80°C.

Storage Buffer:

PBS only, pH7.3


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Non-treated A431 cells and Forskolin treated A431 cells were subjected to SDS PAGE followed by western blot with 80035-2-RR (Phospho-Beta Catenin (Ser552) antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with Beta Actin (66009-1-Ig) antibody as a loading control. This data was developed using the same antibody clone with 80035-2-PBS in a different storage buffer formulation.



Non-treated HeLa cells and Forskolin treated HeLa cells were subjected to SDS PAGE followed by western blot with 80035-2-RR (Phospho-Beta Catenin (Ser552) antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with Beta Actin (66009-1-Ig) antibody as a loading control. This data was developed using the same antibody clone with 80035-2-PBS in a different storage buffer formulation.