

For Research Use Only

# Phospho-GYS (Ser641) Recombinant antibody, PBS Only

Catalog Number: 80102-2-PBS



## Basic Information

<b>Catalog Number:</b> 80102-2-PBS	<b>GenBank Accession Number:</b> BC007688	<b>Purification Method:</b> Protein A purification
<b>Size:</b> 100ug, Concentration: 1 mg/ml by Nanodrop;	<b>GeneID (NCBI):</b> 2997	<b>CloneNo.:</b> 250744A11
<b>Source:</b> Rabbit	<b>UNIPROT ID:</b> P13807	
<b>Isotype:</b> IgG	<b>Full Name:</b> glycogen synthase 1 (muscle)	
	<b>Calculated MW:</b> 84 kDa	
	<b>Observed MW:</b> 84 kDa	

## Applications

**Tested Applications:**  
WB, Indirect ELISA  
**Species Specificity:**  
human, mouse

## Background Information

Glycogen synthase 1 (GYS1, GS) catalyzes the key step of glycogen synthesis and plays an important role in glycogen metabolism in the liver and muscle. In kidney tissues, glycogen synthase 1 (GYS1) is the most important rate-limiting enzyme functioning in the last step of glycogen synthesis. Pathologically, its deficiency has been shown to cause muscle glycogen storage disease type 0 and death. Studies of tumors showed that GYS1 was rapidly induced under hypoxic conditions and positively correlated with glycogen accumulation in glioblastoma, breast, and colon cancer cell lines. GYS1 is phosphorylated at nine sites, and insulin stimulates the dephosphorylation of glycogen synthase. Insulin stimulates dephosphorylation of glycogen synthase via PKB-mediated phosphorylation of GSK3. Phosphorylation of GSK3 decreases kinase activity, which will decrease phosphorylation of GS and increase glycogen synthase fractional activity. (PMID: 32802186, PMID: 30443599, PMID: 22232606)


## Storage

**Storage:**  
Store at -80°C.  
**Storage Buffer:**  
PBS only, pH7.3

For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Non-treated NIH/3T3 cells, phosphatase inhibitor treated and λ phosphatase NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 80102-2-RR (Phospho-GYS (Ser641) antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with Alpha Tubulin antibody (66031-1-Ig) as loading control. This data was developed using the same antibody clone with 80102-2-PBS in a different storage buffer formulation.

