For Research Use Only

PARP1 Recombinant antibody

Catalog Number:80174-1-RR

1 Publications



Purification Method:

Recommended Dilutions:

WB: Jurkat cells, HeLa cells, ROS1728 cells, HepG2

IHC: human liver cancer tissue, mouse testis tissue

IF: mouse testis tissue, MCF-7 cells, HepG2 cells

Basic Information

Catalog Number: GenBank Accession Number:

80174-1-RR BC037545 Protein A purification GeneID (NCBI): CloneNo.:

100ul, Concentration: 250 µg/ml by 142 3N19 Nanodrop;

Source: poly (ADP-ribose) polymerase 1

WB 1:5000-1:20000 Rabbit IHC 1:50-1:500 Calculated MW: IF 1:50-1:500 Isotype: 1014 aa, 113 kDa

IgG Observed MW: Immunogen Catalog Number: 113-116, 89 kDa

AG4193

Applications

Tested Applications:

IF, IHC, WB, ELISA **Cited Applications:**

Species Specificity: Human, mouse, rat **Cited Species:**

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

PARP1 (poly(ADP-ribose) polymerase 1) is a nuclear enzyme catalyzing the poly(ADP-ribosyl)ation of many key proteins in vivo. The normal function of PARP1 is the routine repair of DNA damage. Activated by DNA strand breaks, the PARP1 is cleaved into an 85 to 89-kDa COOH-terminal fragment and a 24-kDa NH2-terminal peptide by caspases during the apoptotic process. The appearance of PARP fragments is commonly considered as an important biomarker of apoptosis. In addition to caspases, other proteases like calpains, cathepsins, granzymes and matrix metalloproteinases (MMPs) have also been reported to cleave PARP1 and gave rise to fragments ranging from 42-89-kDa. This antibody was generated against the C-terminal region of human PARP1 and it recognizes the fulllength as well as the cleavage of the PARP1.

Positive Controls:

cells, RAW264.7 cells

Notable Publications

Author **Pubmed ID** Journal Application WB Yue Liu Int J Biol Macromol 35533849

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Jurkat cells treated with Staurosporin were subjected to SDS PAGE followed by western blot with 80174-1-RR (PARP1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5




Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 80174-1-RR (PARP1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (4% PFA) fixed mouse testis tissue using PARP1 antibody (80174-1-RR, Clone: 3N19) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed mouse testis tissue using PARP1 antibody (80174-1-RR, Clone: 3N19) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 80174-1-RR (PARP1 antibody) at dilution of 1:2000 (under 40x Lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using PARP1 antibody (80174-1-RR, Clone: 3N19) at dilution of 1:100 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).