

For Research Use Only

Phospho-AMPK Alpha (Thr172) Recombinant monoclonal antibody, PBS Only

Catalog Number: 80209-6-PBS



Basic Information

Catalog Number: 80209-6-PBS	GenBank Accession Number: BC048980	Purification Method: Protein A purification
Size: 100ug , Concentration: 1 mg/ml by Nanodrop;	GeneID (NCBI): 5562	CloneNo.: 242923D4
Source: Rabbit	UNIPROT ID: Q13131	
Isotype: IgG	Full Name: protein kinase, AMP-activated, alpha 1 catalytic subunit	
	Observed MW: 64 kDa	

Applications

Tested Applications:
WB, Indirect ELISA

Species Specificity:
human, mouse

Background Information

AMPK is a serine/threonine protein kinase complex consisting of a catalytic α -subunit ($\alpha 1$ and $\alpha 2$), a scaffolding β -subunit ($\beta 1$ and $\beta 2$), and a regulatory γ -subunit ($\gamma 1$, $\gamma 2$, and $\gamma 3$). Ubiquitous expression of AMPK $\alpha 1$ -, $\beta 1$ -, and $\gamma 1$ -subunits in many tissues makes the $\alpha 1\beta 1\gamma 1$ complex a reference for AMPK assays to identify AMPK activators. AMPK is generally quiescent under normal conditions but is activated in response to signals and stresses that increase the AMP/ATP ratio, such as hypoglycemia, strenuous exercise, anoxia, and ischemia. An increase in the ratio of AMP/ATP activates AMPK by several mechanisms, including direct allosteric activation and covalent modification in which an AMP-dependent AMPK kinase (AMPKK) phosphorylates the α subunit on Thr172. Once activated, AMPK switches on catabolic pathways that generate ATP, while switching off ATP-consuming processes (e.g., biosynthesis, cell growth, and proliferation), and in doing so acts as an "energy gauge". (PMID: 27034026, PMID: 21980456, PMID: 27600021)

This antibody can recognize the phosphorylation sites of Thr183 in AMPK Alpha 1 and Thr172 in AMPK Alpha 2.

Storage


Storage:
Store at -80°C.

Storage Buffer:
PBS only, pH7.3


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Non-treated C2C12 cells and H2O2 treated C2C12 cells were subjected to SDS PAGE followed by western blot with 80209-6-RR (Phospho-AMPK Alpha (Thr172) antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with GAPDH (60004-1-Ig) antibody as a loading control. This data was developed using the same antibody clone with 80209-6-PBS in a different storage buffer formulation.



Non-treated, phosphatase inhibitor treated and λ phosphatase treated HEK-293 cells were subjected to SDS PAGE followed by western blot with 80209-6-RR (Phospho-AMPK Alpha (Thr172) antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with Beta Actin (66009-1-Ig) antibody as a loading control. This data was developed using the same antibody clone with 80209-6-PBS in a different storage

