

For Research Use Only

Di/Tri-Methyl-Histone H3 (Lys9) Recombinant monoclonal antibody, PBS Only



Catalog Number: 80240-2-PBS

Basic Information

Catalog Number: 80240-2-PBS	GenBank Accession Number: BC066245	Purification Method: Protein A purification
Size: 100ug , Concentration: 1 mg/ml by Nanodrop;	GeneID (NCBI): 8350	CloneNo.: 241994C6
Source: Rabbit	UNIPROT ID: P68431	
Isotype: IgG	Full Name: histone cluster 1, H3a	
	Observed MW: 15 kDa	

Applications

Tested Applications:
WB, IF/ICC, Dot Blot, Indirect ELISA, ChIP-qPCR

Species Specificity:
human, mouse

Background Information

Histones, including H1/H5 (linker histones), H2, H3, and H4 (core histones), are nucleic proteins which interact with DNA to form the nucleosomes and play important roles in gene regulation and DNA replication. Histone proteins are highly post-translationally modified while Histone H3 is the most extensively modified. Methylation of Histone H3 at lysine 9 is linked to transcriptional repression.

Storage


Storage:
Store at -80°C.

Storage Buffer:
PBS only, pH7.3


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Dot blot analysis was used to confirm the specificity of 80240-2-RR Di/Tri-Methyl-Histone H3 (Lys9) antibody. peptides were spotted onto NC and probed with antibody at 1 µg/mL. The amount of peptide (µg/mL) spotted is indicated next to each row. This data was developed using the same antibody clone with 80240-2-PBS in a different storage buffer formulation.




Chromatin was prepared from HeLa cells. Cells were fixed with formaldehyde for 10 minutes. The ChIP was performed with 15 µg of cross-linked chromatin, 5 µg of Di/Tri-Methyl-Histone H3 (Lys9) (80240-2-RR) or 5 µg of Normal Rabbit IgG (98136-1-RR), and 20 µl of Protein A Magarose Beads. The immunoprecipitated DNA was quantified by real-time PCR. This data was developed using the same antibody clone with 80240-2-PBS in a different storage buffer formulation.



NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 80240-2-RR (HIST1H3A antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 80240-2-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 80240-2-RR (Di/Tri-Methyl-Histone H3 (Lys9) antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 80240-2-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed NIH/3T3 cells using Di/Tri-Methyl-Histone H3 (Lys9) antibody (80240-2-RR, Clone: 241994C6) at dilution of 1:1000 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). This data was developed using the same antibody clone with 80240-2-PBS in a different storage buffer formulation.