For Research Use Only

## P62,SQSTM1 Recombinant antibody, **PBS Only**

Catalog Number:80294-1-PBS

**Featured Product** 



**Purification Method:** 

Protein A purification

CloneNo.:

1D17

**Basic Information** 

Catalog Number: GenBank Accession Number: BC017222

80294-1-PBS GeneID (NCBI): Size:

100ug, Concentration: 1mg/ml by

Nanodrop: **UNIPROT ID:** Q13501 Rabbit Full Name: Isotype: sequestosome 1 IgG Calculated MW:

Immunogen Catalog Number: 48 kDa AG13131

Observed MW:

62 kDa

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, FC (Intra), IP, Indirect ELISA

Species Specificity:

human


**Background Information** 


Sequestosome 1 (SQSTM1/p62) is a multifunctional adaptor protein implicated in selective autophagy, cell signaling pathways, and tumorigenesis. p62 has been implicated in shuttling ubiquitinated and aggregated proteins for autophagic degradation. p62 is degraded during the autophagic process, which makes its intracellular level a  $marker for autophagy\ progression.\ p62\ is\ at\ the\ cross-roads\ of\ several\ signaling\ pathways\ including\ Ras/\ Raf/\ MAPK$ and NFkB and plays important role in cancer. p62 is a component of inclusion bodies/ protein aggregates found in human diseases, including Huntington's disease, Alzheimer's disease, Parkinson's disease, and nephropathic cystinosis (PMID: 22074114, 22860231, 22714671).


Storage

Storage: Store at -80°C. Storage Buffer: PBS Only

## Selected Validation Data










Lysates of HepG2 cells treated with Chloroquine or not were subjected to SDS PAGE followed by western blot with 80294-1-RR (P62,SQSTM1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 80294-1-PBS in a different storage buffer formulation.

WB result of P62,SQSTM1 antibody (80294-1-RR; 1:600; incubated at room temperature for 1.5 hours) with sh-Control and sh-P62,SQSTM1 transfected HEK-293 cells. This data was developed using the same antibody clone with 80294-1-PBS in a different storage buffer formulation.

Various lysates were subjected to SDS PAGE followed by western blot with 80294-1-RR (P62,SQSTM1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 80294-1-PBS in a different storage buffer formulation.








Immunofluorescent analysis of (-20°C Ethanol) fixed Chloroquine treated HeLa cells using P62,SQSTM1 antibody (80294-1-RR, Clone: 1D17) at dilution of 1:500 and Coralite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red). This data was developed using the same antibody clone with 80294-1-PBS in a different storage buffer formulation.

1X10^6 Jurkat cells were intracellularly stained with 0.4 ug Anti-Human P62,5Q5TM1 (80294-1-RR, Clone:1D17) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 80294-1-PBS in a different storage buffer formulation.

IP result of anti-P62,SQSTM1 (IP:80294-1-RR, 4ug; Detection:80294-1-RR 1:500) with HEK-293 cells lysate 1640 ug. This data was developed using the same antibody clone with 80294-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human pancreas cancer tissue slide using 80294-1-RR (P62,SQSTM1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 80294-1-PBS in a different storage buffer formulation.