For Research Use Only

Phospho-eEF2K (Ser366) Recombinant antibody

Catalog Number:80351-3-RR



Purification Method:

Basic Information

Catalog Number: GenBank Accession Number:

80351-3-RR BC032665 Protein A purification

GeneID (NCBI): CloneNo.: 100ul , Concentration: 1000 $\mu g/ml$ by 29904 250361E10

Nanodrop: **UNIPROT ID:** Recommended Dilutions: Source: 000418 WB: 1:500-1:2000

Rabbit Full Name:

Isotype: eukaryotic elongation factor-2 kinase

IgG Calculated MW:

725 aa, 100 kDa Observed MW: 100 kDa

Applications

Tested Applications:

WB, ELISA

Species Specificity:

human

Positive Controls:

WB: Jurkat cells,

Background Information

Eukaryotic Elongation Factor-2 Kinase (eEF2K) acts as a negative regulator of protein synthesis, translation, and cell growth. As a structurally unique member of the alpha-kinase family, eEF2K is essential to cell survival under stressful conditions, as it contributes to both cell viability and proliferation. eEF2K is regulated by various $mechanisms, including phosphorylation through residues and autophosphorylation. \\ee F2K is regulated by various and autophosphorylation for the residues and autophosphorylation for the residues and autophosphorylation. \\$ mechanisms, including phosphorylation through residues and autophosphorylation. eEF2K is downregulated $through \ the \ phosphory lation \ of \ multiple \ sites \ via \ mTOR \ signaling \ and \ upregulated \ via \ the \ AMPK \ pathway. \ In \ the$ S6K-mediated pathway, RPS6KB1 or p70S6K, can phosphorylate eEF2K on Ser-366, rendering it inactive (PMID: 11500364, PMID: 34532346). The calculated molecular weight of eEF2K is 82kDa, and probably due to phosphorylation modification, resulting in a larger molecular weight, around 100kDa.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


*** 20ul sizes contain 0.1% BSA

in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Jurkat cell lysates were subjected to SDS PAGE followed by western blot with 80351-3-RR (Phospho-eEF2K (Ser366) antibody) blocked with BSA only, Phospho-eEF2K (Ser366) peptide or non-Phospho peptide at dilution of 1:1000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with Lamin B1 (66095-1-lg) antibody as a loading control.