

For Research Use Only

Phospho-JNK (Thr183/Tyr185) Recombinant antibody, PBS Only

Catalog Number: 80435-3-PBS



Basic Information

Catalog Number:	80435-3-PBS	GenBank Accession Number:	NM_138982	Purification Method:	Protein A purification
Size:	100ug, Concentration: 1 mg/ml by Nanodrop;	GeneID (NCBI):	5599	CloneNo.:	243163F8
Source:	Rabbit	UNIPROT ID:	P45983		
Isotype:	IgG	Full Name:	mitogen-activated protein kinase 8		
		Calculated MW:	48 kDa		
		Observed MW:	46 kDa, 54 kDa		

Applications

Tested Applications:

WB, Indirect ELISA

Species Specificity:

human, mouse

Background Information

MAPK8(Mitogen-activated protein kinase 8) is also named as JNK1, PRKM8, SAPK1, SAPK1C and belongs to the MAP kinase subfamily. The JNK gene generates 10 forms of JNK through alternative splicing, and the protein encoded by the JNK gene has or does not have a COOH terminal, resulting in 46 kDa and 54 kDa proteins. MAPK8 is activated by dual phosphorylation at a Thr-Pro-Tyr motif during response to UV light. Phosphorylation of these sites in response to UV results in transcriptional activation of c-Jun. The antibody can detect endogenous levels of p46 and p54 SAPK/JNK when phosphorylated at Thr183 and Tyr185. It will also react with JNK singly phosphorylated at Thr183.

Storage

Storage:

Store at -80°C.

Storage Buffer:

PBS only, pH7.3

For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech
Group brand and is not available to purchase from any
other manufacturer.

Selected Validation Data



Non-treated NIH/3T3 cells and UV treated NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 80435-3-RR (Phospho-JNK (Thr183/Tyr185) antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with Beta Actin (66009-1-Ig) antibody as a loading control. This data was developed using the same antibody clone with 80435-3-PBS in a different storage buffer formulation.

Non-treated HEK-293 cells and UV treated HEK-293 cells were subjected to SDS PAGE followed by western blot with 80435-3-RR (Phospho-JNK (Thr183/Tyr185) antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with Beta Actin (66009-1-Ig) antibody as a loading control. This data was developed using the same antibody clone with 80435-3-PBS in a different storage buffer formulation.