## For Research Use Only

## C1QBP Recombinant antibody

Catalog Number:80490-1-RR



**Basic Information** 

Catalog Number: GenBank Accession Number:

80490-1-RR BC013731 GeneID (NCBI): CloneNo.: 3K20

100ul, Concentration: 1000 µg/ml by 708

Source:

complement component 1, q Rabbit subcomponent binding protein Isotype: Calculated MW:

282 aa, 31 kDa IgG Immunogen Catalog Number: Observed MW: AG19773 32 kDa

**Purification Method:** Protein A purification

Recommended Dilutions:

WB 1:2000-1:10000 IHC 1:50-1:500 IF 1:250-1:1000

**Applications** 

**Tested Applications:** 

FC, IF, IHC, WB, ELISA Species Specificity: Human, Mouse, Rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

**Positive Controls:** 

WB: HeLa cells, HEK-293 cells, HepG2 cells, Jurkat cells, K-562 cells, HSC-T6 cells, PC-12 cells, NIH/3T3 cells, 4T1 cells

IHC: human tonsillitis tissue,

IF: HeLa cells.

## **Background Information**

C1QBP, also named as gC1q receptor (gC1qR), p32, p33, and hyaluronan-binding protein 1 (HABP1), is a protein initially copurified with splicing factor SF2 (PMID: 1830244). The protein is synthesized as a pro-protein of 282 amino acids (aa) that is post-translationally processed by removal of the initial 73 aa to a mature protein of 209 aa (PMID: 8262387). C1QBP is an evolutionary conserved and ubiquitously expressed multifunctional protein and has been reported to be a predominantly mitochondrial matrix protein involved in inflammation and infection processes, mitochondrial ribosome biogenesis, regulation of apoptosis and nuclear transcription, and pre-mRNA splicing (PMID: 28942965).

Storage


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 80490-1-RR (C1QBP antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 80490-1-RR (C1QBP antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 80490-1-RR (C1QBP antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed HeLa cells using C1QBP antibody (80490-1-RR, Clone: 3K20) at dilution of 1:500 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CoraLite® 594 Beta Actin antibody (CL594-66009, Clone: 2D4H5, red).



1X10^6 HeLa cells were intracellularly stained with 0.5 ug Anti-Human C1QBP (80490-1-RR, Clone:3K20) (red) labeled with FlexAble CoraLite® Plus 555 Antibody Labeling Kit for Rabbit IgG (KFA002), or 0.5 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).