For Research Use Only

iNOS Recombinant antibody


Catalog Number:80517-1-RR 43 Publications



Basic Information	Catalog Number: 80517-1-RR	GenBank Accession Number: NM_000625	Purification Method: Protein A purification	
	Size:	GenelD (NCBI):	CloneNo.:	
	100ul , Concentration: 1000 µg/ml by		6022	
	Nanodrop;	UNIPROT ID:	Recommended Dilutions:	
	Source:	P35228	WB 1:2000-1:10000	
	Rabbit	Full Name:	IHC 1:50-1:500	
	Isotype: nitric oxide synthase 2, inducible		e	
	lgG	Calculated MW: 131 kDa		
		Observed MW: 110-130 kDa		
Applications	Tested Applications:	Positive Controls:		
			/ 264.7 cells, LPS treated human PBMCs	
	Cited Applications: WB, IHC, IF		IHC : mouse brain tissue, human colon tissue, rat brain tissue, human liver tissue, human lung tissue	
	Species Specificity: human, mouse, rat			
	Cited Species: human, mouse, rat			
	Note-IHC: suggested antigen r TE buffer pH 9.0; (*) Alternativ retrieval may be performed w	vely, antigen		
	buffer pH 6.0			
Background Information	NOS2, also named as iNOS and NOS2 functions throughout the body. NO is including neurotransmission, antimic expressed in liver and is inducible by short half-life due to rapid degradation	a reactive free radical which acts robial and antitumoral activities. a combination of lipopolysaccha on by calpain. iNOS monomer is a CaM-binding site(503-532aa) of i	ch is a messenger molecule with diverse as a biologic mediator in several processes, NOS2 is a nitric oxide synthase which is ride and certain cytokines. iNOS has a very direct substrate of calpain I and can be NOS, and then a ~70kD band can be detected	
	NOS2, also named as iNOS and NOS2 functions throughout the body. NO is including neurotransmission, antimic expressed in liver and is inducible by short half-life due to rapid degradatic cleaved by calpain I at the canonical by western (PMID:11786228). This an	a reactive free radical which acts robial and antitumoral activities. a combination of lipopolysaccha on by calpain. iNOS monomer is a CaM-binding site(503-532aa) of i tibody is specific to NOS2.	as a biologic mediator in several processes, NOS2 is a nitric oxide synthase which is ride and certain cytokines. iNOS has a very direct substrate of calpain I and can be NOS, and then a ~70kD band can be detected	
	NOS2, also named as iNOS and NOS2 functions throughout the body. NO is including neurotransmission, antimic expressed in liver and is inducible by short half-life due to rapid degradatic cleaved by calpain I at the canonical by western (PMID:11786228). This an Author Pub	a reactive free radical which acts robial and antitumoral activities. a combination of lipopolysaccha on by calpain. iNOS monomer is a CaM-binding site(503-532aa) of i tibody is specific to NOS2. med ID Journal	as a biologic mediator in several processes, NOS2 is a nitric oxide synthase which is ride and certain cytokines. iNOS has a very direct substrate of calpain I and can be NOS, and then a ~70kD band can be detected Application	
	NOS2, also named as iNOS and NOS2 functions throughout the body. NO is including neurotransmission, antimic expressed in liver and is inducible by short half-life due to rapid degradatic cleaved by calpain I at the canonical by western (PMID:11786228). This an Author Pub Songya Li 360	a reactive free radical which acts robial and antitumoral activities. a combination of lipopolysaccha on by calpain. iNOS monomer is a CaM-binding site(503-532aa) of i tibody is specific to NOS2. med ID Journal 76540 Biomed Pharmac	as a biologic mediator in several processes, NOS2 is a nitric oxide synthase which is ride and certain cytokines. iNOS has a very direct substrate of calpain I and can be NOS, and then a ~70kD band can be detected Application other IHC	
	NOS2, also named as iNOS and NOS2 functions throughout the body. NO is including neurotransmission, antimic expressed in liver and is inducible by short half-life due to rapid degradatic cleaved by calpain I at the canonical by western (PMID:11786228). This an Author Pub Songya Li 360 Tianchuan Zhu 397	a reactive free radical which acts robial and antitumoral activities. a combination of lipopolysaccha on by calpain. iNOS monomer is a CaM-binding site(503-532aa) of i tibody is specific to NOS2. med ID Journal 76540 Biomed Pharmac	as a biologic mediator in several processes, NOS2 is a nitric oxide synthase which is ride and certain cytokines. iNOS has a very direct substrate of calpain I and can be NOS, and then a ~70kD band can be detected Application other IHC IHC	
	NOS2, also named as iNOS and NOS2 functions throughout the body. NO is including neurotransmission, antimic expressed in liver and is inducible by short half-life due to rapid degradatic cleaved by calpain I at the canonical by western (PMID:11786228). This an Author Pub Songya Li 360 Tianchuan Zhu 397	a reactive free radical which acts robial and antitumoral activities. a combination of lipopolysaccha on by calpain. iNOS monomer is a CaM-binding site(503-532aa) of i tibody is specific to NOS2. med ID Journal 76540 Biomed Pharmac	as a biologic mediator in several processes, NOS2 is a nitric oxide synthase which is ride and certain cytokines. iNOS has a very direct substrate of calpain I and can be NOS, and then a ~70kD band can be detected Application other IHC IHC	
Notable Publications	NOS2, also named as iNOS and NOS2 functions throughout the body. NO is- including neurotransmission, antimic expressed in liver and is inducible by short half-life due to rapid degradatic cleaved by calpain I at the canonical by western (PMID:11786228). This an Author Pub Songya Li 360 Tianchuan Zhu 397 Yang Yu 395 Storage: Store at -20°C. Stable for one year aft Storage Buffer: PBS with 0.02% sodium azide and 50	a reactive free radical which acts robial and antitumoral activities. a combination of lipopolysaccha on by calpain. iNOS monomer is a CaM-binding site(503-532aa) of i tibody is specific to NOS2. med ID Journal 76540 Biomed Pharmac 47173 Nat Commun 11926 Cell Biochem Fun er shipment. % glycerol, pH7.3	as a biologic mediator in several processes, NOS2 is a nitric oxide synthase which is ride and certain cytokines. iNOS has a very direct substrate of calpain I and can be NOS, and then a ~70kD band can be detected Application other IHC IHC	
Background Information Notable Publications Storage *** 20ul sizes contain 0.1% BSA	NOS2, also named as iNOS and NOS2 functions throughout the body. NO is- including neurotransmission, antimic expressed in liver and is inducible by short half-life due to rapid degradatic cleaved by calpain I at the canonical by western (PMID:11786228). This an Author Pub Songya Li 360 Tianchuan Zhu 397 Yang Yu 395 Storage: Store at -20°C. Stable for one year aft Storage Buffer:	a reactive free radical which acts robial and antitumoral activities. a combination of lipopolysaccha on by calpain. iNOS monomer is a CaM-binding site(503-532aa) of i tibody is specific to NOS2. med ID Journal 76540 Biomed Pharmac 47173 Nat Commun 11926 Cell Biochem Fun er shipment. % glycerol, pH7.3	as a biologic mediator in several processes, NOS2 is a nitric oxide synthase which is ride and certain cytokines. iNOS has a very direct substrate of calpain I and can be NOS, and then a ~70kD band can be detected Application other IHC IHC	


T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data




Vivue Pret AB colu

EDTA buffer (pH 9.0).




Untreated and LPS treated RAW 264.7 cells were subjected to SDS PAGE followed by western blot with 80517-1-RR (iNOS antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.


Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 80517-1-RR (iNOS antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 80517-1-RR (iNOS antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded rat brain tissue slide using 80517-1-RR (iNOS antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human colon tissue slide using 80517-1-RR (iNOS antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).





Immunohistochemical analysis of paraffinembedded human colon tissue slide using 80517-1-RR (iNOS antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

Immunohistochemical analysis of paraffinembedded rat brain tissue slide using 80517-1-RR (iNOS antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).