For Research Use Only

## NRF2, NFE2L2 Recombinant antibody, PBS Only



Catalog Number:80593-1-PBS

**Featured Product** 

**Basic Information** 

Catalog Number:

GenBank Accession Number: BC011558

**Purification Method:** 

80593-1-PBS

GeneID (NCBI):

Protein A purification

100ug, Concentration: 1 mg/ml by

CloneNo.: 1|21

Nanodrop:

**UNIPROT ID:** Q16236 Full Name:

Rabbit Isotype:

nuclear factor (erythroid-derived 2)-

IgG

like 2

Immunogen Catalog Number: AG9489

Calculated MW: 605 aa. 68 kDa

Observed MW:

110 kDa

**Applications** 

**Tested Applications:** 

WB, IP, IF, FC, IHC, Indirect ELISA

Species Specificity:

Human, mouse


## **Background Information**

NRF2, also named as NFE2L2, belongs to the bZIP family and CNC subfamily. It is a transcription activator that binds to antioxidant response (ARE) elements in the promoter regions of target genes. NRF2 is important for the coordinated up-regulation of genes in response to oxidative stress. It may be involved in the transcriptional activation of genes of the beta-globin cluster by mediating enhancer activity of hypersensitive site 2 of the betaglobin locus control region. Nrf2 is a key player in the regulation of genes encoding for many antioxidative response enzymes.The expression of NRF2 may be induced under oxidative stress (PMID:14567983).In lung cancer, Nrf2 activation in malignant cells has been associated with tumor progression and chemotherapy resistance(PMID:20534738). Identifying patients with abnormal NRF2 expression may be important for selection for chemotherapy in NSCLC. As new investigators break into the emerging field of Nrf2 research, confusion regarding the correct migratory pattern of Nrf2 is causing doubts about the accuracy and reproducibility of published results. This letter provides solid evidence that the actually observed molecular weight of Nrf2 is about 70kDa and 95-110 kDa. (PMID: 22703241).


Storage

Storage: Store at -80°C. Storage Buffer: PBS Only


## **Selected Validation Data**



Untreated and MG132 treated HEK-293 cells were subjected to SDS PAGE followed by western blot with 80593-1-RR (NRF2, NFE2L2 antibody) at dilution of 1:2500 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 80593-1-PBS in a different storage buffer formulation.




Untreated and MG132 treated HepG2 cells were subjected to SDS PAGE followed by western blot with 80593-1-RR (NRF2, NFE2L2 antibody) at dilution of 1:2500 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 80593-1-PBS in a different storage buffer formulation.




Untreated and MG132 treated HeLa cells were subjected to SDS PAGE followed by western blot with 80593-1-RR (NRF2, NFE2L2 antibody) at dilution of 1:2500 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 80593-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 80593-1-RR (NRF2, NFE2L2 antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 80593-1-PBS in a different storage buffer formulation.




1X10^6 MCF-7 cells were intracellularly stained with 0.4 ug Anti-Human NRF2, NFE2L2 (80593-1-RR, Clone:1121) and Coralite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011). This data was developed using the same antibody clone with 80593-1-PBS in a different storage buffer formulation.




IP result of anti-NRF2, NFE2L2 (IP:80593-1-RR, 4ug; Detection:80593-1-RR 1:700) with HeLa cells lysate 1520 ug. This data was developed using the same antibody clone with 80593-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 80593-1-RR (NRF2, NFE2L2 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 80593-1-PBS in a different storage buffer formulation




Immunohistochemical analysis of paraffinembedded human pancreas cancer tissue slide using 80593-1-RR (NRF2, NFE2L2 antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 80593-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human pancreas cancer tissue slide using 80593-1-RR (NRF2, NFE2L2 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 80593-1-PBS in a different storage buffer formulation.





Immunofluorescent analysis of (-20°C Ethanol) fixed MG132 treated HepG2 cells using NRF2, NFE2L2 antibody (80593-1-RR, Clone: 1121) at dilution of 1:600 and Coralite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red). This data was developed using the same antibody clone with 80593-1-PBS in a different storage buffer formulation.

Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using NRF2, NFE2L2 antibody (80593-1-RR, Clone: 1121) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). This data was developed using the same antibody clone with 80593-1-PBS in a different storage buffer formulation.